arXiv:1410.0170v1 [math.DG] 1 Oct 2014

Multiply Warped Products with a Quarter-symmetric Connection

Quan Qu, Yong Wang *

Abstract

In this paper, we study the Einstein warped products and multiply warped products with a quarter-symmetric connection. We also study warped products and multiply warped products with a quarter-symmetric connection with constant scalar curvature. Then apply our results to generalized Robertson-Walker space-times with a quarter-symmetric connection and generalized Kasner space-times with a quarter-symmetric connection.

Keywords: Warped products; multiply warped products; quarter-symmetric connection; Ricci tensor; scalar curvature; Einstein manifolds.

1 Introduction

The (singly) warped product $B \times_b F$ of two pseudo-Riemannian manifolds (B, g_B) and (F, g_F) with a smooth function $b : B \to (0, \infty)$ is a product manifold of form $B \times F$ with the metric tensor $g = g_B \oplus b^2 g_F$. Here, (B, g_B) is called the base manifold and (F, g_F) is called as the fiber manifold and b is called as the warping function. The concept of warped products was first introduced by Bishop and O'Neill [1] to construct examples of Riemannian manifolds with negative curvature. In [2], F. Dobarro and E. Dozo had studied the problem of showing when a Riemannian metric of constant scalar curvature can be produced on a product manifolds by a warped product construction from the viewpoint of partial differential equations and variational methods. In [3],

*Corresponding author.

E-mail addresses: quq453@nenu.edu.cn (Q. Qu), wangy581@nenu.edu.cn (Y. Wang).

School of Mathematics and Statistics, Northeast Normal University, Changchun Jilin, 130024, China

Ehrlich, Jung and Kim got explicit solutions to warping function to have a constant scalar curvature for generalized Robertson-Walker space-times. In [4], explicit solutions were also obtained for the warping function to make the space-time as Einstein when the fiber is also Einstein.

One can generalize (singly) warped products to multiply warped products. A multiply warped product (M, g) is the product manifold $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$ with the metric $g = g_B \oplus b_1^2 g_{F_1} \oplus b_2^2 g_{F_2} \cdots \oplus b_m^2 g_{F_m}$, where for each $i \in \{1, \cdots, m\}, b_i : B \to (0, \infty)$ is smooth and (F_i, g_{F_i}) is a pseudo-Riemannian manifold. In particular, when $B = (c, d)$, the metric $g_B = -dt^2$ is negative and (F_i, g_{F_i}) is a Riemannian manifold, we call M as the multiply generalized Robertson-Walker space-time.

Singly warped products have a natural generalization. A twisted product (M, g) is a product manifold of form $M = B \times_b F$, with a smooth function $b : B \times F \to (0, \infty)$, and the metric tensor $g = g_B \oplus b^2 g_F$. In [5], they showed that mixed Ricci-flat twisted products could be expressed as warped products. As a consequence, any Einstein twisted products are warped products. Similar to the definition of multiply warped product, a multiply twisted product (M, g) is a product manifold of form $M = B \times b_1 F_1 \times b_2 F_2 \cdots \times b_m F_m$ with the metric $g = g_B \oplus b_1^2 g_{F_1} \oplus b_2^2 g_{F_2} \cdots \oplus b_m^2 g_{F_m}$, where for each $i \in \{1, \cdots, m\}, b_i : B \times F_i \to (0, \infty)$ is smooth. So in this paper, we define the multiply twisted products as generalizations of twisted products and multiply warped products.

The definition of a semi-symmetric metric connection was given by H. Hayden in [6]. In 1970, K. Yano [7] considered a semi-symmetric metric connection and studied some of its properties. Then in 1975, Golab [8] introduced the idea of a quarter-symmetric linear connection in differentiable manifold which is a generalization of semi-symmetric connection. A linear connection ∇ on an n-dimensional Riemannian manifold (M, g) is called a quarter-symmetric connection if its torsion tensor T of the connection ∇ satisfies $T(X, Y) = \pi(Y)\phi X - \pi(X)\phi Y$, where π is a 1-form and ϕ is a (1,1) tensor field. In particular, if $\phi(X) = X$, then the quarter-symmetric connection reduces to the semi-symmetric connection.

In [9], Dobarro and Ünal studied Ricci-flat and Einstein-Lorentzian multiply warped products and considered the case of having constant scalar curvature for multiply warped products and applied their results to generalized Kasner space-times. In [10], S. Sular and C. Özgür studied warped product manifolds with a semi-symmetric metric connection, they computed curvature of semi-symmetric metric connection and consid-

ered Einstein warped product manifolds with a semi-symmetric metric connection. In [11], they studied warped product manifolds with a semi-symmetric non-metric connection. In [12], we considered multiply warped products with a semi-symmetric metric connection, then apply our results to generalized Robertson-Walker spacetimes with a semi-symmetric metric connection and generalized Kasner spacetimes with a semi-symmetric metric connection. In [13], we studied curvature of multiply warped products with a semi-symmetric non-metric connection. In this paper, we will generalize our result to warped and multiply warped products with a quarter-symmetric connection.

This paper is arranged as follows. In Section 2, we give the definition of a quarter-symmetric connection and its curvature, then give the formula of the Levi-Civita connection and curvature of singly warped and multiply twisted product. In section 3, we first compute curvature of a singly warped product with a quarter-symmetric connection, then study the generalized Robertson-Walker space-times with a quarter-symmetric connection. In section 4, firstly we compute curvature of multiply twisted products with a quarter-symmetric connection, secondly we study the special multiply warped product with a quarter-symmetric connection, finally we consider the generalized Kasner space-times with a quarter-symmetric connection.

2 Preliminaries

Let M be a Riemannian manifold with Riemannian metric g. A linear connection $\overline{\nabla}$ on a Riemannian manifold M is called a quarter-symmetric connection if the torsion tensor T of the connection $\overline{\nabla}$

$$T(X,Y) = \overline{\nabla}_X Y - \overline{\nabla}_Y X - [X,Y] \tag{1}$$

satisfies

$$T(X,Y) = \pi(Y)\phi X - \pi(X)\phi Y \tag{2}$$

where π is a 1-form associated with the vector field P on M defined by $\pi(X) = g(X,P)$ and ϕ is a (1,1) tensor field. $\overline{\nabla}$ is called a quarter-symmetric metric connection if it satisfies $\overline{\nabla}g = 0$. $\overline{\nabla}$ is called a quarter-symmetric non-metric connection if it satisfies $\overline{\nabla}g \neq 0$.

If ∇ is the Levi-Civita connection of M, in the equation (2.4) in [14], let $\varphi_1 = \lambda_1,\ \varphi_2 = 0,\ U = P,\ f_1 = 0,\ f_2 = \lambda_2 - \lambda_1,\ U_2 = P,\ \lambda_1 \neq 0,\ \lambda_2 \neq 0$, then we get a

linear connection $\overline{\nabla}$ defined by

$$\overline{\nabla}_X Y = \nabla_X Y + \lambda_1 \pi(Y)X - \lambda_2 g(X,Y)P. \tag{3}$$

It is easy to see that:
when $\lambda_1 = \lambda_2 = 1$, $\overline{\nabla}$ is a semi-symmetric metric connection;
when $\lambda_1 = \lambda_2 \neq 1$, $\overline{\nabla}$ is a quarter-symmetric metric connection;
when $\lambda_1 \neq \lambda_2$, $\overline{\nabla}$ is a quarter-symmetric non-metric connection.

In [12], we considered the case . In this paper, we will consider cases and .

Let R and $\overline{R}$ be the curvature tensors of ∇ and $\overline{\nabla}$, respectively. By the equation (3.13) in [14], we can get

$$\begin{aligned} \overline{R}(X,Y)Z &= R(X,Y)Z + \lambda_1 g(Z,\nabla_X P)Y - \lambda_1 g(Z,\nabla_Y P)X \\ &+ \lambda_2 g(X,Z)\nabla_Y P - \lambda_2 g(Y,Z)\nabla_X P \\ &+ \lambda_1\lambda_2 \pi(P)[g(X,Z)Y - g(Y,Z)X] \\ &+ \lambda_2^2[g(Y,Z)\pi(X) - g(X,Z)\pi(Y)]P \\ &+ \lambda_1^2 \pi(Z)[\pi(Y)X - \pi(X)Y] \end{aligned} \tag{4}$$

for any vector fields X, Y, Z on M.

Remark 1. *When $\lambda_1 = \lambda_2 = 1$, we can get the equation* (4) *in* [12].

2.1 Warped Product

Let (B, g_B) and (F, g_F) be two Riemannian manifolds and $f : B \to (0, \infty)$ be a smooth function. The warped product is the product manifold $B \times F$ with the metric tensor $g = g_B \oplus f^2 g_F$. The function f is called the warping function of the warped product, and the Hessian of f is defined by $H^f(X,Y) = XYf - (\nabla_X Y)f$.

We need the following two lemmas from [14], for later use:

Lemma 2.1. *Let $M = B \times_f F$ be a warped product, ∇, ∇^B and ∇^F denote the Levi-Civita connection on M, B and F, respectively. If $X, Y \in \Gamma(TB)$ and $U, W \in \Gamma(TF)$, then:*
(1)$\nabla_X Y = \nabla^B_X Y$;
(2)$\nabla_X U = \nabla_U X = \frac{Xf}{f} U$;
(3)$\nabla_U W = -\frac{g(U,W)}{f} grad_B f + \nabla^F_U W$.

Lemma 2.2. *Let $M = B \times_f F$ be a warped product with curvature R, If $X, Y, Z \in \Gamma(TB)$ and $U, V, W \in \Gamma(TF)$, then:*
$(1) R(X,Y)Z = R^B(X,Y)Z$;
$(2) R(V,X)Y = -\frac{H^f_B(X,Y)}{f}V$;
$(3) R(X,Y)V = R(V,W)X = 0$;
$(4) R(X,V)W = -\frac{g(V,W)}{f}\nabla^B_X grad_B f$;
$(5) R(V,W)U = R^F(V,W)U - \frac{|grad_B f|^2_B}{f^2}[g(W,U)V - g(V,U)W]$.

2.2 Multiply Twisted Product

A multiply twisted product (M, g) is a product manifold of form $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$ with the metric $g = g_B \oplus b_1^2 g_{F_1} \oplus b_2^2 g_{F_2} \cdots \oplus b_m^2 g_{F_m}$, where for each $i \in \{1, \cdots, m\}, b_i : B \times F_i \to (0, \infty)$ is smooth. Similarly the Hessian of b_i is defined by $H^{b_i}(X,Y) = XYb_i - (\nabla_X Y)b_i$.

We need the following two lemmas from [12], for later use:

Lemma 2.3. *Let $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$ be a multiply twisted product and let $X, Y \in \Gamma(TB)$ and $U \in \Gamma(TF_i), W \in \Gamma(TF_j)$, then:*
$(1) \nabla_X Y = \nabla^B_X Y$;
$(2) \nabla_X U = \nabla_U X = \frac{Xb_i}{b_i}U$;
$(3) \nabla_U W = 0$ *if* $i \neq j$;
$(4) \nabla_U W = U(lnb_i)W + W(lnb_i)U - \frac{g_{F_i}(U,W)}{b_i} grad_{F_i} b_i - b_i g_{F_i}(U,W) grad_B b_i + \nabla^{F_i}_U W$ *if* $i = j$.

Lemma 2.4. *Let $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$ be a multiply twisted product and let $X, Y, Z \in \Gamma(TB)$ and $V \in \Gamma(TF_i), W \in \Gamma(TF_j), U \in \Gamma(TF_k)$, then:*
$(1) R(X,Y)Z = R^B(X,Y)Z$;
$(2) R(V,X)Y = -\frac{H^{b_i}_B(X,Y)}{b_i}V$;
$(3) R(X,V)W = R(V,W)X = R(V,X)W = 0$ *if* $i \neq j$;
$(4) R(X,Y)V = 0$;
$(5) R(V,W)X = VX(lnb_i)W - WX(lnb_i)V$ *if* $i = j$;
$(6) R(V,W)U = 0$ *if* $i = j \neq k$ *or* $i \neq j \neq k$;
$(7) R(U,V)W = -g(V,W)\frac{g_B(grad_B b_i, grad_B b_k)}{b_i b_k}U$ *if* $i \neq j \neq k$;
$(8) R(X,V)W = -\frac{g(V,W)}{b_i}\nabla^B_X(grad_B b_i) + [WX(lnb_i)]V - g_{F_i}(W,V) grad_{F_i} X(lnb_i)$ *if* $i = j$;
$(9) R(V,W)U = g(V,U) grad_B(W(lnb_i)) - g(W,U) grad_B(V(lnb_i)) + R^{F_i}(V,W)U$
$- \frac{|grad_B b_i|^2_B}{b_i^2}[g(W,U)V - g(V,U)W]$ *if* $i = j = k$.

Remark 2. *It is easy to see that Lemmas* 2.1 *and* 2.2 *are Corollaries of Lemma* 2.3 *and* 2.4, *respectively.*

Finally we define the curvature, Ricci curvature and scalar curvature as follows:

$$R(X,Y) = \nabla_X \nabla_Y - \nabla_Y \nabla_X - \nabla_{[X,Y]},$$

$$Ric(X,Y) = \sum_k \varepsilon_k \langle R(X,E_k)Y, E_k\rangle,$$

$$S = \sum_k \varepsilon_k Ric(E_k,E_k)$$

where E_k is an orthonormal base of M with $\langle E_k, E_k\rangle = \varepsilon_k$, $\varepsilon_k = \pm 1$.

3 Warped Product with a Quarter-symmetric Connection

In this section, we firstly compute curvature, Ricci curvature and scalar curvature of singly warped product with a quarter-symmetric connection, then study the generalized Robertson-Walker space-times with a quarter-symmetric connection.

3.1 Connection and Curvature

By Lemma 2.1 and the equation (3), we have the following two Propositions:

Proposition 3.1. *Let $M = B \times_f F$ be a warped product. If $X, Y \in \Gamma(TB)$, $U, W \in \Gamma(TF)$ and $P \in \Gamma(TB)$, then:*
$(1)\overline{\nabla}_X Y = \overline{\nabla}^B_X Y$;
$(2)\overline{\nabla}_X U = \frac{Xf}{f} U$;
$(3)\overline{\nabla}_U X = \left[\frac{Xf}{f} + \lambda_1 \pi(X)\right] U$;
$(4)\overline{\nabla}_U W = -f g_F(U,W) grad_B f + \nabla^F_U W - \lambda_2 g(U,W) P$.

Proposition 3.2. *Let $M = B \times_f F$ be a warped product. If $X, Y \in \Gamma(TB)$, $U, W \in \Gamma(TF)$ and $P \in \Gamma(TF)$, then:*
$(1)\overline{\nabla}_X Y = \nabla^B_X Y - \lambda_2 g(X,Y) P$;
$(2)\overline{\nabla}_X U = \frac{Xf}{f} U + \lambda_1 \pi(U) X$;
$(3)\overline{\nabla}_U X = \frac{Xf}{f} U$;
$(4)\overline{\nabla}_U W = -\frac{g(U,W)}{f} grad_B f + \overline{\nabla}^F_U W$.

By Lemmas 2.12.2 and the equation (4), we have the following two Propositions:

Proposition 3.3. *Let $M = B\times_f F$ be a warped product. If $X, Y, Z \in \Gamma(TB)$, $U, V, W \in \Gamma(TF)$ and $P \in \Gamma(TB)$, then:*
$(1)\overline{R}(X,Y)Z = \overline{R}^B(X,Y)Z$;
$(2)\overline{R}(V,X)Y = -\Big[\frac{H_B^f(X,Y)}{f} + \lambda_2\frac{Pf}{f}g(X,Y) + \lambda_1\lambda_2\pi(P)g(X,Y) + \lambda_1 g(Y,\nabla_X P)$
$- \lambda_1^2\pi(X)\pi(Y)\Big]V$;
$(3)\overline{R}(X,Y)V = 0$;
$(4)\overline{R}(V,W)X = 0$;
$(5)\overline{R}(X,V)W = -g(V,W)\Big[\frac{\nabla_X^B(grad_B f)}{f} + \lambda_1\frac{Pf}{f}X + \lambda_2\nabla_X P + \lambda_1\lambda_2\pi(P)X - \lambda_2^2\pi(X)P\Big]$;
$(6)\overline{R}(U,V)W = R^F(U,V)W - \Big[\frac{|grad_B f|_B^2}{f^2} + (\lambda_1+\lambda_2)\frac{Pf}{f} + \lambda_1\lambda_2\pi(P)\Big]\Big[g(V,W)U$
$- g(U,W)V\Big]$.

Proposition 3.4. *Let $M = B\times_f F$ be a warped product. If $X, Y, Z \in \Gamma(TB)$, $U, V, W \in \Gamma(TF)$ and $P \in \Gamma(TF)$, then:*
$(1)\overline{R}(X,Y)Z = R^B(X,Y)Z + \lambda_2\Big[g(X,Z)\frac{Yf}{f} - g(Y,Z)\frac{Xf}{f}\Big]P + \lambda_1\lambda_2\pi(P)[g(X,Z)Y$
$- g(Y,Z)X]$;
$(2)\overline{R}(V,X)Y = -\frac{H_B^f(X,Y)}{f}V - \lambda_1\pi(V)\frac{Yf}{f}X - \lambda_2 g(X,Y)\nabla_V P - g(X,Y)[\lambda_1\lambda_2\pi(P)V$
$- \lambda_2^2\pi(V)P]$;
$(3)\overline{R}(X,Y)V = \lambda_1\pi(V)\Big[\frac{Xf}{f}Y - \frac{Yf}{f}X\Big]$;
$(4)\overline{R}(V,W)X = \lambda_1\frac{Xf}{f}[\pi(W)V - \pi(V)W]$;
$(5)\overline{R}(X,V)W = -g(V,W)\frac{\nabla_X^B(grad_B f)}{f} + \lambda_1\frac{Xf}{f}\pi(W)V - \lambda_1 g(W,\nabla_V P)X - \lambda_2 g(V,W)\frac{Xf}{f}P$
$- \lambda_1\lambda_2 g(V,W)\pi(P)X + \lambda_1^2\pi(W)\pi(V)X$;
$(6)\overline{R}(U,V)W = R^F(U,V)W - \frac{|grad_B f|_B^2}{f^2}[g(V,W)U - g(U,W)V] + \lambda_1[g(W,\nabla_U P)V$
$- g(W,\nabla_V P)U] + \lambda_2[g(U,W)\nabla_V P - g(V,W)\nabla_U P)] + \lambda_1\lambda_2\pi(P)[g(U,W)V$
$- g(V,W)U] + \lambda_2^2[g(V,W)\pi(U) - g(U,W)\pi(V)]P + \lambda_1^2\pi(W)[\pi(V)U - \pi(U)V]$.

By Propositions 3.3 and 3.4 and the definition of the Ricci curvature tensor, we have the following two Propositions:

Proposition 3.5. *Let $M = B\times_f F$ be a warped product,* $\dim B = n_1, \dim F = n_2$ *and* $\dim M = \overline{n} = n_1 + n_2$. *If $X, Y \in \Gamma(TB)$, $V, W \in \Gamma(TF)$ and $P \in \Gamma(TB)$, then:*
$(1)\overline{Ric}(X,Y) = \overline{Ric}^B(X,Y) + n_2\Big[\frac{H_B^f(X,Y)}{f} + \lambda_2\frac{Pf}{f}g(X,Y) + \lambda_1\lambda_2\pi(P)g(X,Y)$
$+ \lambda_1 g(Y,\nabla_X P) - \lambda_1^2\pi(X)\pi(Y)\Big]$;
$(2)\overline{Ric}(X,V) = \overline{Ric}(V,X) = 0$;

$(3)\overline{Ric}(V,W) = \overline{Ric}^F(V,W) + \Big\{\frac{\Delta_B f}{f} + (n_2-1)\frac{|grad_B f|_B^2}{f^2} + [(\overline{n}-1)\lambda_1\lambda_2 - \lambda_2^2]\pi(P)$
$+ \lambda_2 div_B P + [(\overline{n}-1)\lambda_1 + (n_2-1)\lambda_2]\frac{Pf}{f}\Big\}g(V,W).$

where $div_B P = \sum_{k=1}^{n_1} \varepsilon_k \langle \nabla_{E_k} P, E_k\rangle$, *and* $E_k, 1 \le k \le n_1$ *is an orthonormal base of* B *with* $\varepsilon_k = g(E_k, E_k)$.

Proposition 3.6. *Let* $M = B \times_f F$ *be a warped product,* $\dim B = n_1, \dim F = n_2$ *and* $\dim M = \overline{n} = n_1 + n_2$. *If* $X, Y \in \Gamma(TB)$, $V, W \in \Gamma(TF)$ *and* $P \in \Gamma(TF)$, *then:*
$(1)\overline{Ric}(X,Y) = \overline{Ric}^B(X,Y) + n_2\frac{H_B^f(X,Y)}{f} + [(\overline{n}-1)\lambda_1\lambda_2 - \lambda_2^2]\pi(P)g(X,Y)$
$+ \lambda_2 g(X,Y) div_F P;$
$(2)\overline{Ric}(X,V) = [(\overline{n}-1)\lambda_1 - \lambda_2]\pi(V)\frac{Xf}{f};$
$(3)\overline{Ric}(V,X) = [\lambda_2 - (\overline{n}-1)\lambda_1]\pi(V)\frac{Xf}{f};$
$(4)\overline{Ric}(V,W) = Ric^F(V,W) + g(V,W)\Big\{\frac{\Delta_B f}{f} + (n_2-1)\frac{|grad_B f|_B^2}{f^2} + [(\overline{n}-1)\lambda_1\lambda_2 - \lambda_2^2]\pi(P)\Big\}$
$+ [(\overline{n}-1)\lambda_1 - \lambda_2]g(W, \nabla_V P) + [\lambda_2^2 + (1-\overline{n})\lambda_1^2]\pi(V)\pi(W) + \lambda_2 g(V,W) div_F P.$

By Proposition 3.5 and the definition of the scalar curvature, we have the following:

Proposition 3.7. *Let* $M = B \times_f F$ *be a warped product,* $\dim B = n_1, \dim F = n_2$ *and* $\dim M = \overline{n} = n_1 + n_2$. *If* $P \in \Gamma(TB)$, *then the scalar curvature* $\overline{S}$ *has the following expression:*

$$\overline{S} = \overline{S}^B + 2n_2\frac{\Delta_B f}{f} + \frac{S^F}{f^2} + n_2(n_2-1)\frac{|grad_B f|_B^2}{f^2} + n_2(\overline{n}-1)(\lambda_1+\lambda_2)\frac{Pf}{f}$$
$$+ [n_2(\overline{n}+n_1-1)\lambda_1\lambda_2 - n_2(\lambda_1^2+\lambda_2^2)]\pi(P) + n_2(\lambda_1+\lambda_2) div_B P.$$

By Proposition 3.6 and the definition of the scalar curvature, we have the following:

Proposition 3.8. *Let* $M = B \times_f F$ *be a warped product,* $\dim B = n_1, \dim F = n_2$ *and* $\dim M = \overline{n} = n_1 + n_2$. *If* $P \in \Gamma(TF)$, *then the scalar curvature* $\overline{S}$ *has the following expression:*

$$\overline{S} = \overline{S}^B + 2n_2\frac{\Delta_B f}{f} + \frac{S^F}{f^2} + n_2(n_2-1)\frac{|grad_B f|_B^2}{f^2} + [\overline{n}(\overline{n}-1)\lambda_1\lambda_2 + (1-\overline{n})(\lambda_1^2+\lambda_2^2)]\pi(P)$$
$$+ (\overline{n}-1)(\lambda_1+\lambda_2) div_F P.$$

3.2 Generalized Robertson-Walker Space-times with a Quarter-symmetric Connection

Theorem 3.9. *Let* $M = I \times_f F$ *be a warped product, where* I *is an open interval in* $\mathbb{R}$, $\dim I = 1$ *and* $\dim F = \overline{n} - 1 (\overline{n} \ge 3)$. *Then* $(M, \overline{\nabla})$ *is Einstein if and only if* (F, ∇^F) *is Einstein for* $P = \frac{\partial}{\partial t}$ *or* f *is a constant on* I *for* $P \in \Gamma(TF)$, $\lambda_2 \ne (\overline{n}-1)\lambda_1$.

Proof. Assume that $P = \frac{\partial}{\partial t}$, Let $f = e^{\frac{q}{2}}$, by Proposition 3.5, we can write

$$\overline{Ric}\Big(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\Big) = (1-\overline{n})\Big[\frac{1}{4}(q')^2 + \frac{1}{2}q'' - \frac{1}{2}\lambda_2 q' + \lambda_1\lambda_2 - \lambda_1^2\Big] g_I\Big(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\Big); \tag{5}$$

$$\overline{Ric}\Big(\frac{\partial}{\partial t}, V\Big) = 0;$$

$$\begin{aligned}\overline{Ric}(V,W) = Ric^F(V,W) + e^q\Big\{\frac{\overline{n}-1}{4}(q')^2 + \frac{1}{2}\Big[(\overline{n}-1)\lambda_1 + (\overline{n}-2)\lambda_2\Big]q' \\ +\lambda_2^2 + (1-\overline{n})\lambda_1\lambda_2 + \frac{1}{2}q''\Big\} g_F(V,W)\end{aligned} \tag{6}$$

for any $V, W \in \Gamma(TF)$.

Since M is Einstein manifold, we have

$$\overline{Ric}\Big(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\Big) = \alpha g_I\Big(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\Big); \tag{7}$$

$$\overline{Ric}(V,W) = \alpha e^q g_F(V,W). \tag{8}$$

From equations (5) and (7), we get

$$\alpha = (1-\overline{n})\Big[\frac{1}{4}(q')^2 + \frac{1}{2}q'' - \frac{1}{2}\lambda_2 q' + \lambda_1\lambda_2 - \lambda_1^2\Big]. \tag{9}$$

Similarly, from equations (6) and (8), and by the use of (9), we obtain

$$Ric^F(V,W) = \Big[\frac{1-\overline{n}}{2}(q')^2 - \frac{\overline{n}}{2}q'' + \Big(\frac{1-\overline{n}}{2}\lambda_1 + \frac{1}{2}\lambda_2\Big)q' + (\overline{n}-1)\lambda_1^2 - \lambda_2^2\Big] e^q g_F(V,W)$$

which implies that (F, ∇^F) is Einstein manifold.

Assume that $P \in \Gamma(TF)$, by Proposition 3.6, we have

$$\overline{Ric}\Big(\frac{\partial}{\partial t}, V\Big) = \frac{1}{2}q'[(\overline{n}-1)\lambda_1 - \lambda_2]\pi(V); \tag{10}$$

$$\overline{Ric}\Big(V, \frac{\partial}{\partial t}\Big) = \frac{1}{2}q'[\lambda_2 - (\overline{n}-1)\lambda_1]\pi(V) \tag{11}$$

for any $V \in \Gamma(TF)$.

Since M is an Einstein manifold, we can write

$$\overline{Ric}\Big(\frac{\partial}{\partial t}, V\Big) = \alpha g\Big(\frac{\partial}{\partial t}, V\Big) = 0 = \alpha g\Big(V, \frac{\partial}{\partial t}\Big) = \overline{Ric}\Big(V, \frac{\partial}{\partial t}\Big) \tag{12}$$

where $\frac{\partial}{\partial t} \in \Gamma(TB)$ and $V \in \Gamma(TF)$.

Since $\lambda_2 \neq (\overline{n}-1)\lambda_1, \pi(V) \neq 0,$, using equations $(10), (11), (12)$, then we can get $q' = 0$, which means q is a constant on I, then f is a constant on I. □

Theorem 3.10. *Let $M = B \times_f I$ be a warped product, where I is an open interval in $\mathbb{R}$, $\dim I = 1$ and $\dim B = \overline{n} - 1$ ($\overline{n} \geq 3$). Then*
(1)If $(M, \overline{\nabla})$ is an Einstein manifold, $P \in \Gamma(TB)$, $\nabla^B P = 0$, and f is a constant on B, then:

$$\overline{S}^B = [(\overline{n}-1)(\overline{n}-2)\lambda_1\lambda_2 + \lambda_1^2 + (1-\overline{n})\lambda_2^2]\pi(P).$$

Furthermore, if $(\overline{n}-1)(\overline{n}-2)\lambda_1\lambda_2 + \lambda_1^2 + (1-\overline{n})\lambda_2^2 = 0$, then $\overline{S}^B = 0$.
(2)If $(M, \overline{\nabla})$ is Einstein manifold, $P \in \Gamma(TI)$, and $\lambda_2 \neq (\overline{n}-1)\lambda_1$ then f is a constant on B.
(3)If f is a constant on B, (B, ∇^B) is Einstein, $P \in \Gamma(TI)$, then $(M, \overline{\nabla})$ is Einstein manifold; furthermore, if $\lambda_2 = (\overline{n}-1)\lambda_1$, then $\alpha = \alpha_B$, where α and α_B denote the Einstein constant on M and B, respectively.

Proof. Assume that $(M, \overline{\nabla})$ is an Einstein manifold, $P \in \Gamma(TB)$, then

$$\overline{Ric}(X,Y) = \frac{\overline{s}}{\overline{n}} g(X,Y) \tag{13}$$

for any $X, Y \in \Gamma(TB)$. Consider that f is a constant on B and by Proposition 3.7, we can get

$$\overline{Ric}(X,Y) = \frac{1}{\overline{n}} g(X,Y)\{\overline{S}^B + [(2\overline{n}-2)\lambda_1\lambda_2 - (\lambda_1^2 + \lambda_2^2)]\pi(P)\}.$$

Define $\widehat{S} = \sum\limits_{k=1}^{\overline{n}-1} \frac{1}{\overline{n}} \overline{Ric}(E_k, E_k)$, where E_k is an orthonormal base of B, then we have

$$\widehat{S} = \frac{\overline{n}-1}{\overline{n}} \Big\{ \overline{S}^B + [(2\overline{n}-2)\lambda_1\lambda_2 - (\lambda_1^2 + \lambda_2^2)]\pi(P) \Big\}. \tag{14}$$

On the other hand, using Proposition 3.5, we can get

$$\overline{Ric}(X,Y) = \overline{Ric}^B(X,Y) + [\lambda_1\lambda_2\pi(P)g(X,Y) - \lambda_1^2\pi(X)\pi(Y)],$$

then we have

$$\widehat{S} = \overline{S}^B + [(\overline{n}-1)\lambda_1\lambda_2 - \lambda_1^2]\pi(P). \tag{15}$$

From equations (14) and (15), we can get

$$\overline{S}^B = [(\overline{n}-1)(\overline{n}-2)\lambda_1\lambda_2 + \lambda_1^2 + (1-\overline{n})\lambda_2^2]\pi(P).$$

Assume that $(M, \overline{\nabla})$ is Einstein manifold, $P \in \Gamma(TI)$, by Proposition 3.6, we obtain

$$\overline{Ric}(X,P) = [(\overline{n}-1)\lambda_1 - \lambda_2]\pi(P)\frac{Xf}{f}; \tag{16}$$

$$\overline{Ric}(P,X) = [\lambda_2 - (\overline{n}-1)\lambda_1]\pi(P)\frac{Xf}{f}. \tag{17}$$

Using the similar proof of Theorem 3.9., we can get $Xf = 0$, which means f is a constant on B.

If f is a constant on B, and (B, ∇^B) is Einstein, and $P \in \Gamma(TI)$, then

$$Ric^B(X,Y) = \alpha_B g(X,Y). \tag{18}$$

By Theorem 3.6, we have

$$\overline{Ric}(X,Y) = \overline{Ric}^B(X,Y) + [(\overline{n}-1)\lambda_1\lambda_2 - \lambda_2^2]\pi(P)g(X,Y). \tag{19}$$

So by equations (18) and (19), we can easily get

$$\overline{Ric}(X,Y) = \{\alpha_B + [(\overline{n}-1)\lambda_1\lambda_2 - \lambda_2^2]\pi(P)\}g(X,Y)$$

which means $(M, \overline{\nabla})$ is an Einstein manifold.

Furthermore, if $\lambda_2 = (\overline{n}-1)\lambda_1$, then $\overline{Ric}(X,Y) = \alpha_B g(X,Y)$, and $\alpha = \alpha_B$. □

Now we specially study $M = I \times F$ with the metric tensor $-dt^2 + f(t)^2 g_F$, I is an open interval in $\mathbb{R}$. Let begin with the following theorem:

Theorem 3.11. *Let $M = I \times F$ with the metric tensor $-dt^2 + f(t)^2 g_F$, $P = \frac{\partial}{\partial t}$, $\dim F = l$. Then $(M, \overline{\nabla})$ is Einstein with the Einstein constant α if and only if the following conditions are satisfied*
(1) *(F, ∇^F) is Einstein with the Einstein constant α_F;*
(2) *$l\Big(\lambda_2\frac{f'}{f} - \frac{f''}{f} + \lambda_1^2 - \lambda_1\lambda_2\Big) = \alpha$;*
(3) *$\alpha_F - ff'' + (1-l)(f')^2 + [\lambda_2^2 - l\lambda_1\lambda_2 - \alpha]f^2 + [l\lambda_1 + (l-1)\lambda_2]ff' = 0$.*

Proof. By Proposition 3.5, we have

$$\begin{aligned}
&\overline{Ric}\Big(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\Big) = -l\Big(\lambda_2\frac{f'}{f} - \frac{f''}{f} + \lambda_1^2 - \lambda_1\lambda_2\Big);\\
&\overline{Ric}\Big(\frac{\partial}{\partial t}, V\Big) = \overline{Ric}\Big(V, \frac{\partial}{\partial t}\Big) = 0;\\
&\overline{Ric}(V,W) = Ric^F(V,W) + g_F(V,W)\{-ff'' - (l-1)f'^2 + (\lambda_2^2 - l\lambda_1\lambda_2)f^2\\
&\qquad\qquad +[l\lambda_1 + (l-1)\lambda_2]ff'\}.
\end{aligned}$$

Then by the Einstein condition, we get Theorem 3.11. □

Remark 3. *When* $\lambda_1 = \lambda_2 = 1$, *we can get Corollary* 21 *in* [12].

Considering the dimension of F, we get Corollaries 3.12 and 3.13 of Theorem 3.11:

Corollary 3.12. *Let* $M = I \times F$ *with the metric tensor* $-dt^2 + f(t)^2 g_F, P = \frac{\partial}{\partial t}, \dim F = 1$. *Then* $(M, \overline{\nabla})$ *is Einstein with the Einstein constant* α *if and only if*
$f'' = \lambda_2 f' + (\lambda_1^2 - \lambda_1\lambda_2)f - \alpha f$

Remark 4. (1)*From Theorem* 3.11, *we can also get: if* $\dim F = 1$, *then* $\alpha_F = 0$;

(2) *When* $\lambda_1 = \lambda_2 = 1$, *we can get Corollary* 23 *in* [12].

Corollary 3.13. *Let* $M = I \times F$ *with the metric tensor* $-dt^2 + f(t)^2 g_F, P = \frac{\partial}{\partial t}, \dim F = l > 1$. *Then* $(M, \overline{\nabla})$ *is Einstein with the Einstein constant* α *if and only if the following conditions are satisfied:*
(1) (F, ∇^F) *is Einstein with the Einstein constant* α^F;
(2)$f'' = \lambda_2 f' + (\lambda_1^2 - \lambda_1\lambda_2 - \frac{\alpha}{l})f$;
(3) $\frac{\alpha_F}{1-l} + (f')^2 + \left[\frac{\alpha}{l} + \lambda_1\lambda_2 + \frac{\lambda_2^2 - \lambda_1^2}{1-l}\right] f^2 + \left[\frac{l}{1-l}\lambda_1 + \frac{(l-2)}{1-l}\lambda_2\right] f f' = 0.$

Remark 5. *When* $\lambda_1 = \lambda_2 = 1$, *we can get Corollary* 24 *in* [12].

By Corollary 3.12 and elementary methods for ordinary differential equations, we get:

Theorem 3.14. *Let* $M = I \times F$ *with the metric tensor* $-dt^2 + f(t)^2 g_F, P = \frac{\partial}{\partial t}, \dim F = 1$. *Then* $(M, \overline{\nabla})$ *is Einstein with the Einstein constant* α *if and only if*
$(1)\alpha < (\lambda_1 - \frac{1}{2}\lambda_2)^2, f(t) = c_1 e^{((\lambda_2 + \sqrt{(2\lambda_1 - \lambda_2)^2 - 4\alpha})/2)t} + c_2 e^{((\lambda_2 - \sqrt{(2\lambda_1 - \lambda_2)^2 - 4\alpha})/2)t}$,
$(2)\alpha = (\lambda_1 - \frac{1}{2}\lambda_2)^2, f(t) = c_1 e^{(\lambda_2/2)t} + c_2 t e^{(\lambda_2/2)t}$,
$(3)\alpha > (\lambda_1 - \frac{1}{2}\lambda_2)^2, f(t) = c_1 e^{(\lambda_2/2)t} cos((\sqrt{4\alpha - (2\lambda_1 - \lambda_2)^2}/2)t)$
$+ c_2 e^{(\lambda_2/2)t} sin((\sqrt{4\alpha - (2\lambda_1 - \lambda_2)^2}/2)t)$.

Remark 6. *When* $\lambda_1 = \lambda_2 = 1$, *we can get Corollary* 25 *in* [12].

As a Corollary of Theorem 3.14, we have

Corollary 3.15. *Let* $M = I \times F$ *with the metric tensor* $-dt^2 + f(t)^2 g_F, P = \frac{\partial}{\partial t}, \dim F = 1$, *and* $\lambda_2 = 2\lambda_1$, *then* $(M, \overline{\nabla})$ *is Einstein with the Einstein constant* α *if and only if*
$(1)\alpha < 0, f(t) = c_1 e^{(\lambda_1 + \sqrt{-\alpha})t} + c_2 e^{(\lambda_1 - \sqrt{-\alpha})t}$,
$(2)\alpha = 0, f(t) = c_1 e^{\lambda_1 t} + c_2 t e^{\lambda_1 t}$,
$(3)\alpha > 0, f(t) = c_1 e^{\lambda_1 t} cos(\sqrt{\alpha}t) + c_2 e^{\lambda_1 t} sin(\sqrt{\alpha}t)$.

Theorem 3.16. *Let $M = I \times F$ with the metric tensor $-dt^2 + f(t)^2 g_F$, $P = \frac{\partial}{\partial t}$, $\dim F = l > 1$. Then $(M, \overline{\nabla})$ is Einstein with the Einstein constant α if and only if one of the following conditions is satisfied:*
$(1)\alpha = (\lambda_1^2 - \lambda_1\lambda_2)l,\ \alpha_F = c_2^2(l\lambda_1^2 - \lambda_2^2),\ f(t) = c_2$;
$(2)\lambda_1 = \lambda_2,\ \alpha = 0,\ \alpha_F = (l-1)c_2^2\lambda_1^2,\ f(t) = c_1 e^{\lambda_1 t} + c_2$;
$(3)\lambda_2^2 - 2l\lambda_1^2 + l\lambda_1\lambda_2 \neq 0,\ \lambda_2 \neq l\lambda_1,\ \alpha = \frac{(3l^2+l)\lambda_1^2\lambda_2^2 - (l^2+l)\lambda_1\lambda_1^3 - 2l^2\lambda_1^3\lambda_2}{(l\lambda_1 - \lambda_2)^2},\ \alpha_F = 0,$
$f(t) = c_0 e^{(l\lambda_1^2 - \lambda_2^2)t/(l\lambda_1 - \lambda_2)}$;
$(4)\lambda_2^2 - 2l\lambda_1^2 + l\lambda_1\lambda_2 = 0,\ \alpha = l(\lambda_1 - \frac{1}{2}\lambda_2)^2,\ \alpha_F = 0,\ f(t) = c_1 e^{\frac{\lambda_2}{2}t}$.

Proof. Let $\frac{\alpha}{l} = d_0,\ \frac{\alpha_F}{1-l} = \overline{d_0},\ a_0 = \frac{\lambda_2 + \sqrt{(2\lambda_1 - \lambda_2)^2 - 4d_0}}{2},\ b_0 = \frac{\lambda_2 - \sqrt{(2\lambda_1 - \lambda_2)^2 - 4d_0}}{2}$, then $a_0 + b_0 = \lambda_2,\ a_0 b_0 = d_0 + \lambda_1\lambda_2 - \lambda_1^2$.
$d_0 < (\lambda_1 - \frac{1}{2}\lambda_2)^2$, then $f(t) = c_1 e^{a_0 t} + c_2 e^{b_0 t}$. By Corollary 3.13(3), then

$$\begin{aligned}
&\overline{d_0} + c_1^2\Big(a_0^2 + a_0 b_0 + \lambda_1^2 + \frac{\lambda_2^2 - \lambda_1^2}{1-l} + \frac{l}{1-l}\lambda_1 a_0 + \frac{l-2}{1-l}\lambda_2 a_0\Big)e^{2a_0 t} \\
&\quad + c_2^2\Big(b_0^2 + a_0 b_0 + \lambda_1^2 + \frac{\lambda_2^2 - \lambda_1^2}{1-l} + \frac{l}{1-l}\lambda_1 b_0 + \frac{l-2}{1-l}\lambda_2 b_0\Big)e^{2b_0 t} \qquad (20) \\
&\quad + c_1 c_2\Big[4a_0 b_0 + \frac{2\lambda_2^2 - 2l\lambda_1^2}{1-l} + \Big(\frac{l}{1-l}\lambda_1 + \frac{l-2}{1-l}\lambda_2\Big)(a_0 + b_0)\Big]e^{(a_0+b_0)t} = 0.
\end{aligned}$$

$1) b_0 = 0$, we have $d_0 = \lambda_1^2 - \lambda_1\lambda_2 < (\lambda_1 - \frac{1}{2}\lambda_2)^2,\ a_0 = \lambda_2,\ \alpha = (\lambda_1^2 - \lambda_1\lambda_2)l$. By the equation (20), we get

$$\overline{d_0} + c_1^2\frac{l\lambda_1(\lambda_2 - \lambda_1)}{1-l}e^{2\lambda_2 t} + c_2^2\frac{\lambda_2^2 - l\lambda_1^2}{1-l} + c_1 c_2\frac{l(\lambda_2 + 2\lambda_1)(\lambda_2 - \lambda_1)}{1-l}e^{\lambda_2 t} = 0.$$

Since $e^{2\lambda_2 t}$ and $e^{\lambda_2 t}$ are linearly independent, we have

$$\begin{cases} c_1^2\frac{l\lambda_1(\lambda_2 - \lambda_1)}{1-l} = 0 \\ \overline{d_0} + c_2^2\frac{\lambda_2^2 - l\lambda_1^2}{1-l} = 0 \\ c_1 c_2\frac{l(\lambda_2 + 2\lambda_1)(\lambda_2 - \lambda_1)}{1-l} = 0 \end{cases}$$

$1'. c_1 = 0,\ c_2 \neq 0$. We have $\underline{\alpha = (\lambda_1^2 - \lambda_1\lambda_2)l,\ \alpha_F = c_2^2(l\lambda_1^2 - \lambda_2^2),\ f(t) = c_2}$.
$2'. c_1 \neq 0,\ c_2 = 0$. We have $\underline{\lambda_1 = \lambda_2,\ \alpha = \alpha_F = 0,\ f(t) = c_1 e^{\lambda_1 t}}$.
$3'. c_1 \neq 0,\ c_2 \neq 0$. We have $\underline{\lambda_1 = \lambda_2,\ \alpha = 0,\ \alpha_F = (l-1)c_2^2\lambda_1^2,\ f(t) = c_1 e^{\lambda_1 t} + c_2}$.
It is easy to see that the conclusion of $2'$ is a special case of $3'$.

$2) b_0 \neq 0$, $1'$. $c_1 = 0,\ c_2 \neq 0$. Since $e^{2a_0 t}, e^{2b_0 t}$ and $e^{(a_0+b_0)t}$ are linearly independent, we have $\overline{d_0} = 0,\ b_0^2 + a_0 b_0 + \lambda_1^2 + \frac{\lambda_2^2 - \lambda_1^2}{1-l} + \frac{l}{1-l}\lambda_1 b_0 + \frac{l-2}{1-l}\lambda_2 b_0 = 0$, then $\alpha_F = 0,\ \frac{l\lambda_1 - \lambda_2}{1-l}b_0 =$

$\frac{l\lambda_1^2-\lambda_2^2}{1-l}$. If $\lambda_2 = l\lambda_1$, we have $0 = l\lambda_1^2$, this is a contradiction with $\lambda_1 \neq 0$. So $\lambda_2 \neq l\lambda_1$ and $b_0 = \frac{l\lambda_1^2-\lambda_2^2}{l\lambda_1-\lambda_2}$. Considering that $b_0 < \frac{\lambda_2}{2}$, we get $\frac{\lambda_2^2+l\lambda_1\lambda_2-2l\lambda_1^2}{l\lambda_1-\lambda_2} > 0$. $a_0 = \frac{l\lambda_1(\lambda_2-\lambda_1)}{l\lambda_1-\lambda_2}, d_0 = \frac{(3l+1)\lambda_1^2\lambda_2^2-(l+1)\lambda_1\lambda_2^3-2l\lambda_1^3\lambda_2}{(l\lambda_1-\lambda_2)^2}$, $\alpha = \frac{(3l^2+l)\lambda_1^2\lambda_2^2-(l^2+l)\lambda_1\lambda_2^3-2l^2\lambda_1^3\lambda_2}{(l\lambda_1-\lambda_2)^2}$ and d_0 satisfies $d_0 < (\lambda_1 - \frac{1}{2}\lambda_2)^2$. So in this case, we obtain

$\frac{\lambda_2^2+l\lambda_1\lambda_2-2l\lambda_1^2}{l\lambda_1-\lambda_2} > 0,\ \alpha = \frac{(3l^2+l)\lambda_1^2\lambda_2^2-(l^2+l)\lambda_1\lambda_2^3-2l^2\lambda_1^3\lambda_2}{(l\lambda_1-\lambda_2)^2},\ \alpha_F = 0,\ f(t) = c_2 e^{\frac{l\lambda_1^2-\lambda_2^2}{l\lambda_1-\lambda_2}t}.$

$2'.\ c_1 \neq 0,\ c_2 = 0$. Using the same method we can get $a_0 = \frac{l\lambda_1^2-\lambda_2^2}{l\lambda_1-\lambda_2},\ b_0 = \frac{l\lambda_1(\lambda_2-\lambda_1)}{l\lambda_1-\lambda_2}$ and

$\frac{\lambda_2^2+l\lambda_1\lambda_2-2l\lambda_1^2}{l\lambda_1-\lambda_2} < 0,\ \alpha = \frac{(3l^2+l)\lambda_1^2\lambda_2^2-(l^2+l)\lambda_1\lambda_2^3-2l^2\lambda_1^3\lambda_2}{(l\lambda_1-\lambda_2)^2},\ \alpha_F = 0,\ f(t) = c_1 e^{\frac{l\lambda_1^2-\lambda_2^2}{l\lambda_1-\lambda_2}t}.$

So by $1'$ and $2'$, we get Theorem 3.16(3).

$3'.\ c_1 \neq 0,\ c_2 \neq 0$. Since e^{2a_0t}, e^{2b_0t} and $e^{(a_0+b_0)t}$ are linearly independent, we have

$$a_0^2 + a_0b_0 + \lambda_1^2 + \frac{\lambda_2^2-\lambda_1^2}{1-l} + \frac{l}{1-l}\lambda_1 a_0 + \frac{l-2}{1-l}\lambda_2 a_0 = 0; \tag{21a}$$

$$b_0^2 + a_0b_0 + \lambda_1^2 + \frac{\lambda_2^2-\lambda_1^2}{1-l} + \frac{l}{1-l}\lambda_1 b_0 + \frac{l-2}{1-l}\lambda_2 b_0 = 0. \tag{21b}$$

$(21a) - (21b)$ we get $(a_0 - b_0)\lambda_2 + \frac{l\lambda_1}{1-l}(a_0 - b_0) + \frac{l-2}{1-l}\lambda_2(a_0 - b_0) = 0$, since $a_0 \neq b_0$ we have $\lambda_2 + \frac{l\lambda_1}{1-l} + \frac{l-2}{1-l}\lambda_2 = 0$, then $\lambda_2 = l\lambda_1$, using $(21a)$ again we get $0 = l\lambda_1^2$, this is a contradiction with $\lambda_1 \neq 0$. So in case $3'$ we have no solution.

$d_0 = (\lambda_1 - \frac{1}{2}\lambda_2)^2$, then $f(t) = c_1 e^{\frac{\lambda_2}{2}t} + c_2 t e^{\frac{\lambda_2}{2}t}$. By Corollary 3.13(3), we get

$$\begin{aligned}&\overline{d_0} + \Big(\frac{\lambda_2}{2}c_1 + c_2 + \frac{\lambda_2}{2}c_2 t\Big)^2 e^{\lambda_2 t} + \frac{(5-l)\lambda_2^2 - 4l\lambda_1^2}{4(1-l)}(c_1 + c_2 t)^2 e^{\lambda_2 t}\\ &+\Big(\frac{l}{1-l}\lambda_1 + \frac{l-2}{1-l}\lambda_2\Big)(c_1 + c_2 t)\Big(\frac{\lambda_2}{2}c_1 + c_2 + \frac{\lambda_2}{2}c_2 t\Big)e^{\lambda_2 t} = 0\end{aligned}$$

$1'.\ c_1 = 0,\ c_2 \neq 0$. The coefficient of $e^{\lambda_2 t}$ is $c_2^2 = 0$, this is a contradiction with $c_2 \neq 0$.

$2'.\ c_1 \neq 0,\ c_2 = 0$. Then $\overline{d_0} = 0$ and from the coefficient of $e^{\lambda_2 t}$ we can get $\lambda_2^2 + l\lambda_1\lambda_2 - 2l\lambda_1^2 = 0$, so $\alpha = d_0 l = l(\lambda_1 - \frac{1}{2}\lambda_2)^2,\ \alpha_F = 0,\ f(t) = c_1 e^{\frac{\lambda_2}{2}t}$.

$3'.\ c_1 \neq 0,\ c_2 \neq 0$. From the coefficient of $t^2 e^{\lambda_2 t}$ we get

$$\lambda_2^2 + l\lambda_1\lambda_2 - 2l\lambda_1^2 = 0, \tag{22}$$

by the equation (22), the coefficient of $te^{\lambda_2 t}$ becomes

$$\frac{(l-3)\lambda_2 + 2l\lambda_1}{2(1-l)}c_2 - \frac{\lambda_2^2}{4}c_1 = 0, \tag{23}$$

the coefficient of $e^{\lambda_2 t}$ is $\frac{l\lambda_1-\lambda_2}{1-l}c_1c_2 + c_2^2 = 0$, then $c_2 = \frac{\lambda_2-l\lambda_1}{1-l}c_1$, so by equation (23) we get $(2l-7-l^2)\lambda_2 = (4l^2-10l)\lambda_1$, considering that $2l-7-l^2 \neq 0, 4l^2-10l \neq 0$, we obtain

$\lambda_2 = \frac{(4l^2-10l)}{(2l-7-l^2)}\lambda_1$, using the equation (22) again, we have $(l-1)^2(3l^2-15l+49)=0$, but $l-1 \neq 0, 3l^2-15l+49 \neq 0$, so we have no solution in this case.
$d_0 > (\lambda_1 - \frac{1}{2}\lambda_2)^2$, let $h_0 = \frac{1}{2}\sqrt{4d_0-(2\lambda_1-\lambda_2)^2}$, then $f(t) = e^{\frac{\lambda_2}{2}t}(c_1 cos(h_0 t) + c_2 sin(h_0 t))$. By Corollary 3.13(3), we get

$$\overline{d_0} + \Big[\Big(\frac{\lambda_2}{2}c_1 + c_2 h_0\Big)cos(h_0 t) + \Big(\frac{\lambda_2}{2}c_2 - c_1 h_0\Big)sin(h_0 t)\Big]^2 e^{\lambda_2 t}$$
$$+\Big(d_0 + \lambda_1\lambda_2 + \frac{\lambda_2^2-\lambda_1^2}{1-l}\Big)\Big(c_1 cos(h_0 t) + c_2 sin(h_0 t)\Big)^2 e^{\lambda_2 t} + \Big(\frac{l}{1-l}\lambda_1 + \frac{l-2}{1-l}\lambda_2\Big)$$
$$\Big(c_1 cos(h_0 t) + c_2 sin(h_0 t)\Big)\Big[\Big(\frac{\lambda_2}{2}c_1 + c_2 h_0\Big)cos(h_0 t) + \Big(\frac{\lambda_2}{2}c_2 - c_1 h_0\Big)sin(h_0 t)\Big]e^{\lambda_2 t} = 0.$$

The coefficient of $cos^2(h_0 t)e^{\lambda_2 t}$ is

$$\Big[d_0 + \frac{(l+1)\lambda_2^2 + (4-2l)\lambda_1\lambda_2 - 4\lambda_1^2}{4(1-l)}\Big]c_1^2 + c_2^2 h_0^2 - \frac{l\lambda_1-\lambda_2}{1-l}c_1 c_2 h_0 = 0. \tag{24}$$

Similarly the coefficient of $sin^2(h_0 t)e^{\lambda_2 t}$ is

$$\Big[d_0 + \frac{(l+1)\lambda_2^2 + (4-2l)\lambda_1\lambda_2 - 4\lambda_1^2}{4(1-l)}\Big]c_2^2 + c_1^2 h_0^2 - \frac{l\lambda_1-\lambda_2}{1-l}c_1 c_2 h_0 = 0. \tag{25}$$

If $c_1 = 0,\ c_2 \neq 0$, by the equation (24) we get $c_2^2 h_0^2 = 0$, then $h_0 = 0$, this is a contradiction with $h_0 \neq 0$;
If $c_1 \neq 0,\ c_2 = 0$, by the equation (25) we get $c_1^2 h_0^2 = 0$, then $h_0 = 0$, this is a contradiction with $h_0 \neq 0$. So $c_1 \neq 0,\ c_2 \neq 0$, (24) + (25) we get

$$d_0 = \frac{[(2l-4)\lambda_1 - l\lambda_2](\lambda_2-\lambda_1)}{4(1-l)}. \tag{26}$$

Considering $d_0 > (\lambda_1 - \frac{1}{2}\lambda_2)^2$, we have

$$\lambda_2^2 - 2l\lambda_1^2 + l\lambda_1\lambda_2 > 0. \tag{27}$$

The coefficient of $sin(h_0 t)cos(h_0 t)e^{\lambda_2 t}$ is

$$\frac{\lambda_2^2 - 2l\lambda_1^2 + l\lambda_1\lambda_2}{1-l}c_1 c_2 + \frac{l\lambda_1-\lambda_2}{1-l}h_0(c_2^2 - c_1^2) = 0. \tag{28}$$

Using (24) − (25), we get

$$\frac{\lambda_2^2 - 2l\lambda_1^2 + l\lambda_1\lambda_2}{2(1-l)}(c_1^2 - c_2^2) + \frac{2(l\lambda_1-\lambda_2)}{1-l}h_0 c_1 c_2. \tag{29}$$

Let $a = \frac{\lambda_2^2 - 2l\lambda_1^2 + l\lambda_1\lambda_2}{2(1-l)} < 0,\ b = \frac{l\lambda_1 - \lambda_2}{1-l}h_0,\ x = c_1^2 - c_2^2,\ y = c_1c_2 \neq 0$, from equations (28) and (29) we obtain

$$ax + 2by = 0, \tag{30a}$$

$$2ay - bx = 0. \tag{30b}$$

Through $(30a) \times b + (30b) \times a$, then $a^2 + b^2 = 0$, which means $\left[\frac{\lambda_2^2 - 2l\lambda_1^2 + l\lambda_1\lambda_2}{2(1-l)}\right]^2 + \left[\frac{l\lambda_1 - \lambda_2}{1-l}h_0\right]^2 = 0$, then we get

$$d_0 = \frac{(3l+1)\lambda_1^2\lambda_2^2 - (l+1)\lambda_1\lambda_2^3 - 2l\lambda_1^3\lambda_2}{(l\lambda_1 - \lambda_2)^2}. \tag{31}$$

Using equations (26) and (31), we have $(\lambda_1 - \lambda_2)[(2-l)\lambda_1 - \lambda_2](\lambda_2^2 + l\lambda_1\lambda_2 - 2l\lambda_2^2) = 0$. By inequality (27), we get $\lambda_1 = \lambda_2$ or $\lambda_2 = (2-l)\lambda_1$.

1)$\lambda_1 = \lambda_2$, by the equation (26) we get $d_0 = 0$, this is a contradiction with $d_0 > (\lambda_1 - \frac{1}{2}\lambda_2)^2 = \frac{1}{4}\lambda_1^2 > 0$.

2)$\lambda_2 = (2-l)\lambda_1$, by the equation (26) we get $d_0 = \frac{l^2-4}{4}\lambda_1^2 < \frac{l^2}{4}\lambda_1^2$. On the other hand, $d_0 > (\lambda_1 - \frac{1}{2}\lambda_2)^2 = \frac{l^2}{4}\lambda_1^2$, this is a contradiction.

In a word, we have no solution in the case of $d_0 > (\lambda_1 - \frac{1}{2}\lambda_2)^2$. □

Remark 7. *When $\lambda_1 = \lambda_2 = 1$, we get Theorem 26 in [12].*

Proposition 3.17. *Let $M = I \times F$ with the metric tensor $-dt^2 + f(t)^2 g_F, P = \frac{\partial}{\partial t}$. Then $(M, \overline{\nabla})$ has constant scalar curvature $\overline{S}$ if and only if (F, ∇^F) has constant scalar curvature S^F and*

$$\overline{S} = \frac{S^F}{f^2} - 2l\frac{f''}{f} - l(l-1)\frac{(f')^2}{f^2} + l^2(\lambda_1 + \lambda_2)\frac{f'}{f} + l[\lambda_1^2 + \lambda_2^2 - (l+1)\lambda_1\lambda_2]. \tag{32}$$

Proof. Considering that $M = I \times F$ with the metric tensor $-dt^2 + f(t)^2 g_F$ and $P = \frac{\partial}{\partial t}$, then by Proposition 3.7 we get the equation (32). With the fact that S^F is function defined on F, and f is function defined on I, then using variables separation we complete the proof of this Proposition. □

Proposition 3.18. *Let $M = I \times F$ with the metric tensor $-dt^2 + f(t)^2 g_F,\ P \in \Gamma(TF)$. If $(M, \overline{\nabla})$ has constant scalar curvature $\overline{S}$, then*

(1)*If $\lambda_1 + \lambda_2 \neq 0$ and $\lambda_1^2 + \lambda_2^2 - \overline{n}\lambda_1\lambda_2 = 0$, and $div_F P$ is a constant, then S^F is a constant;*

(2)*If $\lambda_1 = -\lambda_2 \neq 0$ and $g_F(P, P)$ is a constant, then S^F is a constant;*

(3)*If* $\lambda_1+\lambda_2\neq 0$ *and* $\lambda_1^2+\lambda_2^2-\overline{n}\lambda_1\lambda_2\neq 0$, *and* $div_F P$, $g_F(P,P)$ *are constants, then* S^F *is a constant.*

Proof. Considering that $M=I\times F$ with the metric tensor $-dt^2+f(t)^2 g_F$, $P\in\Gamma(TF)$, then by Proposition 3.8, we can get

$$\overline{S}=\frac{S^F}{f^2}-2l\frac{f''}{f}-l(l-1)\frac{(f')^2}{f^2}+[l\overline{n}\lambda_1\lambda_2-l(\lambda_1^2+\lambda_2^2)]f^2 g_F(P,P)+l(\lambda_1+\lambda_2)div_F P. \quad (33)$$

Then by variables separation, we have:
(1)If $\lambda_1+\lambda_2\neq 0$ and $\lambda_1^2+\lambda_2^2-\overline{n}\lambda_1\lambda_2=0$, and $div_F P$ is a constant, we can obtain $\overline{S}=\frac{S^F}{f^2}-2l\frac{f''}{f}-l(l-1)\frac{(f')^2}{f^2}+l(\lambda_1+\lambda_2)div_F P$. Then S^F is a constant;
(2)If $\lambda_1=-\lambda_2\neq 0$ and $g_F(P,P)$ is a constant, which means $\lambda_1+\lambda_2=0$ and $\lambda_1^2+\lambda_2^2-\overline{n}\lambda_1\lambda_2\neq 0$, and $g_F(P,P)$ is a constant, we can obtain $\overline{S}=\frac{S^F}{f^2}-2l\frac{f''}{f}-l(l-1)\frac{(f')^2}{f^2}+[l\overline{n}\lambda_1\lambda_2-l(\lambda_1^2+\lambda_2^2)]f^2 g_F(P,P)$. then S^F is a constant;
(3)It is obvious.
(4)If $\lambda_1+\lambda_2=0$ and $\lambda_1^2+\lambda_2^2-\overline{n}\lambda_1\lambda_2=0$, then we can get $\lambda_1=\lambda_2=0$, which is a contradiction. □

In (32), we make the change of variable $f(t)=\sqrt{v(t)}$ and have the following equation:

$$v''(t)+\frac{l-3}{4}\frac{v'(t)^2}{v(t)}-\frac{l}{2}(\lambda_1+\lambda_2)v'(t)+\Big[(l+1)\lambda_1\lambda_2-\lambda_1^2-\lambda_2^2+\frac{\overline{S}}{l}\Big]v(t)-\frac{S^F}{l}=0. \quad (34)$$

Remark 8. *When* $\lambda_1=\lambda_2=1$, *equations* (32), (33), (34) *respectively become* (20), (21), (22) *in* [12], *then by Proposition* 3.17, *we can get Corollary* 27 *in* [12], *and by Proposition* 3.18.(3), *we can get Corollary* 28 *in* [12].

Theorem 3.19. *Let* $M=I\times F$ *with the metric tensor* $-dt^2+f(t)^2 g_F$, $P=\frac{\partial}{\partial t}$, *and* $\dim F=l=3$. *Then* $(M,\overline{\nabla})$ *has constant scalar curvature* $\overline{S}$ *if and only if* (F,∇^F) *has constant scalar curvature* S^F *and*
(1)$\overline{S}<\frac{27}{16}(\lambda_1+\lambda_2)^2+3\lambda_1^2+3\lambda_2^2-12\lambda_1\lambda_2$ *and* $\overline{S}\neq 3\lambda_1^2+3\lambda_2^2-12\lambda_1\lambda_2$,
$v(t)=c_1 e^{(\left(\frac{3}{2}(\lambda_1+\lambda_2)+\sqrt{\frac{9}{4}(\lambda_1+\lambda_2)^2-\frac{4}{3}\overline{S}+4\lambda_1^2+4\lambda_2^2-16\lambda_1\lambda_2}\right)/2)t}$
$+c_2 e^{(\left(\frac{3}{2}(\lambda_1+\lambda_2)-\sqrt{\frac{9}{4}(\lambda_1+\lambda_2)^2-\frac{4}{3}\overline{S}+4\lambda_1^2+4\lambda_2^2-16\lambda_1\lambda_2}\right)/2)t}+\frac{S^F}{12\lambda_1\lambda_2-3\lambda_1^2-3\lambda_2^2+\overline{S}}$;
(2)$\overline{S}=\frac{27}{16}(\lambda_1+\lambda_2)^2+3\lambda_1^2+3\lambda_2^2-12\lambda_1\lambda_2$,
$v(t)=c_1 e^{\frac{3}{4}(\lambda_1+\lambda_2)t}+c_2 t e^{\frac{3}{4}(\lambda_1+\lambda_2)t}+\frac{S^F}{12\lambda_1\lambda_2-3\lambda_1^2-3\lambda_2^2+\overline{S}}$;
(3)$\overline{S}>\frac{27}{16}(\lambda_1+\lambda_2)^2+3\lambda_1^2+3\lambda_2^2-12\lambda_1\lambda_2$,

$$v(t) = c_1 e^{\frac{3}{4}(\lambda_1+\lambda_2)t} cos((\Big(\sqrt{\frac{4}{3}\overline{S} - 4\lambda_1^2 - 4\lambda_2^2 + 16\lambda_1\lambda_2 - \frac{9}{4}(\lambda_1+\lambda_2)^2}\Big)/2)t)$$
$$+ c_2 e^{\frac{3}{4}(\lambda_1+\lambda_2)t} sin((\Big(\sqrt{\frac{4}{3}\overline{S} - 4\lambda_1^2 - 4\lambda_2^2 + 16\lambda_1\lambda_2 - \frac{9}{4}(\lambda_1+\lambda_2)^2}\Big)/2)t)$$
$$+ \frac{S^F}{12\lambda_1\lambda_2 - 3\lambda_1^2 - 3\lambda_2^2 + \overline{S}};$$

$(4)\overline{S} = 3\lambda_1^2 + 3\lambda_2^2 - 12\lambda_1\lambda_2$, *and* $\lambda_1 + \lambda_2 \neq 0$, $v(t) = c_1 - \frac{2S^F}{9(\lambda_1+\lambda_2)} + c_2 e^{\frac{3}{2}(\lambda_1+\lambda_2)t}$;

$(5)\overline{S} = 3\lambda_1^2 + 3\lambda_2^2 - 12\lambda_1\lambda_2$, *and* $\lambda_1 + \lambda_2 = 0$, $v(t) = \frac{S^F}{6}t^2 + c_1 t + c_2$.

Proof. If $l = 3$, then we have a simple differential equation as follows:

$$v''(t) - \frac{3}{2}(\lambda_1+\lambda_2)v'(t) + (4\lambda_1\lambda_2 - \lambda_1^2 - \lambda_2^2 + \frac{\overline{S}}{3})v(t) - \frac{S^F}{3} = 0. \tag{35}$$

If $\overline{S} \neq 3\lambda_1^2 + 3\lambda_2^2 - 12\lambda_1\lambda_2$, putting $h(t) = (4\lambda_1\lambda_2 - \lambda_1^2 - \lambda_2^2 + \frac{\overline{S}}{3})v(t) - \frac{S^F}{3}$, we get $h''(t) - \frac{3}{2}(\lambda_1+\lambda_2)h'(t) + (4\lambda_1\lambda_2 - \lambda_1^2 - \lambda_2^2 + \frac{\overline{S}}{3})h(t) = 0$. The above solutions $(1)-(3)$ follow directly from elementary methods for ordinary differential equations.
If $\overline{S} = 3\lambda_1^2 + 3\lambda_2^2 - 12\lambda_1\lambda_2$, and $\lambda_1 + \lambda_2 \neq 0$, then $v''(t) - \frac{3}{2}(\lambda_1+\lambda_2)v'(t) - \frac{S^F}{3} = 0$, and we get solution(4).
If $\overline{S} = 3\lambda_1^2 + 3\lambda_2^2 - 12\lambda_1\lambda_2$, and $\lambda_1 + \lambda_2 = 0$, then $v''(t) - \frac{S^F}{3} = 0$, and we get solution(5). □

Remark 9. *When* $\lambda_1 = \lambda_2 = 1$, *we get Theorem* 29 *in* [12].

Theorem 3.20. *Let* $M = I \times F$ *with the metric tensor* $-dt^2 + f(t)^2 g_F, P = \frac{\partial}{\partial t}$, *and* $\dim F = l \neq 3$ *and* $S^F = 0$. *Then* $(M, \overline{\nabla})$ *has constant scalar curvature* $\overline{S}$ *if and only if*

$(1)\overline{S} < \frac{l^3}{4(l+1)}(\lambda_1+\lambda_2)^2 - l[(l+1)\lambda_1\lambda_2 - \lambda_1^2 - \lambda_2^2]$,

$$v(t) = \Big[c_1 e^{(\Big(\frac{1}{2}(\lambda_1+\lambda_2) + \sqrt{\frac{l^2}{4}(\lambda_1+\lambda_2)^2 - (l+1)\Big[(l+1)\lambda_1\lambda_2 - \lambda_1^2 - \lambda_2^2 + \frac{\overline{S}}{l}\Big]}\Big)/2)t}$$
$$+ c_2 e^{(\Big(\frac{1}{2}(\lambda_1+\lambda_2) - \sqrt{\frac{l^2}{4}(\lambda_1+\lambda_2)^2 - (l+1)\Big[(l+1)\lambda_1\lambda_2 - \lambda_1^2 - \lambda_2^2 + \frac{\overline{S}}{l}\Big]}\Big)/2)t}\Big]^{\frac{4}{l+1}};$$

$(2)\overline{S} = \frac{l^3}{4(l+1)}(\lambda_1+\lambda_2)^2 - l[(l+1)\lambda_1\lambda_2 - \lambda_1^2 - \lambda_2^2], v(t) = \Big[c_1 e^{\frac{l}{4}(\lambda_1+\lambda_2)t} + c_2 t e^{\frac{l}{4}(\lambda_1+\lambda_2)t}\Big]^{\frac{4}{l+1}}$;

$(3)\overline{S} > \frac{l^3}{4(l+1)}(\lambda_1+\lambda_2)^2 - l[(l+1)\lambda_1\lambda_2 - \lambda_1^2 - \lambda_2^2]$,

$$v(t) = \Big[c_1 e^{\frac{l}{4}(\lambda_1+\lambda_2)t} cos(\Big(\sqrt{(l+1)\Big[(l+1)\lambda_1\lambda_2 - \lambda_1^2 - \lambda_2^2 + \frac{\overline{S}}{l}\Big] - \frac{l^2}{4}(\lambda_1+\lambda_2)^2}\Big)/2)t$$
$$+ c_2 e^{\frac{l}{4}(\lambda_1+\lambda_2)t} sin(\Big(\sqrt{(l+1)\Big[(l+1)\lambda_1\lambda_2 - \lambda_1^2 - \lambda_2^2 + \frac{\overline{S}}{l}\Big] - \frac{l^2}{4}(\lambda_1+\lambda_2)^2}\Big)/2)t\Big]^{\frac{4}{l+1}}$$

Proof. In this case, (34) is changed into the simpler form

$$v''(t) + \frac{l-3}{4}\frac{v'(t)^2}{v(t)} - \frac{l}{2}(\lambda_1+\lambda_2)v'(t) + \Big[(l+1)\lambda_1\lambda_2 - \lambda_1^2 - \lambda_2^2 + \frac{\overline{S}}{l}\Big]v(t) = 0. \tag{36}$$

Putting $v(t) = w(t)^{\frac{4}{l+1}}$, then $w(t)$ satisfies the equation $w''(t) - \frac{l}{2}(\lambda_1+\lambda_2)w'(t) + \frac{l+1}{4}\big[(l+1)\lambda_1\lambda_2 - \lambda_1^2 - \lambda_2^2 + \frac{\overline{S}}{l}\big]w(t) = 0$, by the elementary methods for ordinary differential equations, we prove the above theorem. □

Remark 10. *When $\lambda_1 = \lambda_2 = 1$, we get Theorem 30 in [12].*

When $\dim F = l \neq 3$, and $S^F \neq 0$, putting $v(t) = w(t)^{\frac{4}{l+1}}$, then $w(t)$ satisfies the following equation :

$$w''(t) - \frac{l}{2}(\lambda_1+\lambda_2)w'(t) + \frac{l+1}{4}\Big[(l+1)\lambda_1\lambda_2 - \lambda_1^2 - \lambda_2^2 + \frac{\overline{S}}{l}\Big]w(t) - \frac{l+1}{4l}S^F w^{1-\frac{4}{l+1}} = 0. \quad (37)$$

4 Multiply Warped Product with a Quarter-symmetric Connection

In this section, firstly we compute curvature of multiply twisted product with a quarter-symmetric connection, secondly we study the special multiply warped product with a quarter-symmetric connection, finally we consider the generalized Kasner space-times with a quarter-symmetric connection.

4.1 Connection and Curvature

By Lemma 2.3 and the equation (3), we have the following two Propositions:

Proposition 4.1. *Let $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$ be a multiply twisted product, If $X, Y \in \Gamma(TB)$, $U \in \Gamma(TF_i), W \in \Gamma(TF_j)$ and $P \in \Gamma(TB)$. Then:*
(1)$\overline{\nabla}_X Y = \overline{\nabla}^B_X Y$;
(2)$\overline{\nabla}_X U = \frac{Xb_i}{b_i}U$;
(3)$\overline{\nabla}_U X = \Big[\frac{Xb_i}{b_i} + \lambda_1\pi(X)\Big]U$;
(4)$\overline{\nabla}_U W = 0$ *if* $i \neq j$;
(5)$\overline{\nabla}_U W = U(lnb_i)W + W(lnb_i)U - \frac{g_{F_i}(U,W)}{b_i}grad_{F_i}b_i - b_i g_{F_i}(U,W)grad_B b_i + \nabla^{F_i}_U W - \lambda_2 g(U,W)P$ *if* $i = j$.

Proposition 4.2. *Let $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$ be a multiply twisted product. If $X, Y \in \Gamma(TB)$, $U \in \Gamma(TF_i), W \in \Gamma(TF_j)$ and $P \in \Gamma(TF_r)$ for a fixed r, then:*
(1)$\overline{\nabla}_X Y = \nabla^B_X Y - \lambda_2 g(X,Y)P$;
(2)$\overline{\nabla}_X U = \frac{Xb_i}{b_i}U + \lambda_1\pi(U)X$;

$(3)\overline{\nabla}_U X = \frac{Xb_i}{b_i}U$;
$(4)\overline{\nabla}_U W = \lambda_1 g(W,P)U$ *if* $i \neq j$;
$(5)\overline{\nabla}_U W = U(lnb_i)W + W(lnb_i)U - \frac{g_{F_i}(U,W)}{b_i}grad_{F_i}b_i - b_i g_{F_i}(U,W)grad_B b_i + \overline{\nabla}^{F_i}_U W$ *if* $i = j$.

By Lemmas 2.3, 2.4 and the equation (4), we have the following two Propositions:

Proposition 4.3. *Let* $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$ *be a multiply twisted product. If* $X, Y, Z \in \Gamma(TB)$, $V \in \Gamma(TF_i), W \in \Gamma(TF_j), U \in \Gamma(TF_k)$ *and* $P \in \Gamma(TB)$, *then:*
$(1)\overline{R}(X,Y)Z = \overline{R}^B(X,Y)Z$;
$(2)\overline{R}(V,X)Y = -\Big[\frac{H^{b_i}_B(X,Y)}{f} + \lambda_2\frac{Pb_i}{b_i}g(X,Y) + \lambda_1\lambda_2\pi(P)g(X,Y) + \lambda_1 g(Y,\nabla_X P)$
$- \lambda_1^2\pi(X)\pi(Y)\Big]V$;
$(3)\overline{R}(X,Y)V = 0$;
$(4)\overline{R}(V,W)X = VX(lnb_i)W - WX(lnb_i)V$ *if* $i = j$;
$(5)\overline{R}(V,W)U = 0$ *if* $i = j \neq k$ *or* $i \neq j \neq k$;
$(6)\overline{R}(X,V)W = \overline{R}(V,W)X = \overline{R}(V,X)W = 0$ *if* $i \neq j$;
$(7)\overline{R}(X,V)W = WX(lnb_i)V - g(V,W)\Big[\frac{\nabla^B_X(grad_B b_i)}{b_i} + grad_{F_i}\frac{X(lnb_i)}{b_i^2} + \lambda_1\frac{Pb_i}{b_i}X + \lambda_2\nabla_X P$
$+ \lambda_1\lambda_2\pi(P)X - \lambda_2^2\pi(X)P\Big]$ *if* $i = j$;
$(8)\overline{R}(U,V)W = -g(V,W)U\Big[\frac{g_B(grad_B b_i, grad_B b_k)}{b_i b_k} + \lambda_1\frac{Pb_i}{b_i} + \lambda_2\frac{Pb_k}{b_k} + \lambda_1\lambda_2\pi(P)\Big]$ *if* $i = j \neq k$;
$(9)\overline{R}(U,V)W = g(U,W)grad_B(V(lnb_i)) - g(V,W)grad_B(U(lnb_i)) + R^{F_i}(U,V)W -$
$\Big[\frac{|grad_B b_i|^2_B}{b_i^2} + (\lambda_1+\lambda_2)\frac{Pb_i}{b_i} + \lambda_1\lambda_2\pi(P)\Big][g(V,W)U - g(U,W)V]$ *if* $i = j = k$.

Proposition 4.4. *Let* $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$ *be a multiply twisted product. If* $X, Y, Z \in \Gamma(TB)$, $V \in \Gamma(TF_i), W \in \Gamma(TF_j), U \in \Gamma(TF_k)$ *and* $P \in \Gamma(TF_r)$ *for a fixed* r, *then:*
$(1)\overline{R}(X,Y)Z = R^B(X,Y)Z + \lambda_2\Big[g(X,Z)\frac{Yb_r}{b_r} - g(Y,Z)\frac{Xb_r}{b_r}\Big]P + \lambda_1\lambda_2\pi(P)[g(X,Z)Y$
$- g(Y,Z)X]$;
$(2)\overline{R}(V,X)Y = -\frac{H^{b_i}_B(X,Y)}{b_i}V - \lambda_1\lambda_2\pi(P)g(X,Y)V$ *if* $i \neq r$;
$(3)\overline{R}(V,X)Y = -\frac{H^{b_i}_B(X,Y)}{b_i} - \lambda_1\pi(V)\frac{Yb_i}{b_i}X - \lambda_2 g(X,Y)\nabla_V P - g(X,Y)[\lambda_1\lambda_2\pi(P)V$
$- \lambda_2^2\pi(V)P]$ *if* $i = r$;
$(4)\overline{R}(X,Y)V = \lambda_1\pi(V)[\frac{Xb_r}{b_r}Y - \frac{Yb_r}{b_r}X]$;
$(5)\overline{R}(V,W)X = -\lambda_1\delta^r_i\frac{Xb_i}{b_i}\pi(V)W + \lambda_1\delta^r_j\frac{Xb_j}{b_j}\pi(W)V$ *if* $i \neq j$;
$(6)\overline{R}(V,W)X = VX(lnb_i)W - WX(lnb_i)V - \lambda_1\delta^r_i\frac{Xb_i}{b_i}[\pi(V)W - \pi(W)V]$ *if* $i = j$;
$(7)\overline{R}(V,W)U = 0$ *if* $i = j \neq k$ *or* $i \neq j \neq k$;
$(8)\overline{R}(X,V)W = \lambda_1\frac{Xb_r}{b_r}\pi(W)V$ *if* $i \neq j$;

$(9)\overline{R}(X,V)W = WX(lnb_i)V - g(V,W)\frac{\nabla^B_X(grad_B b_i)}{b_i} - grad_{F_i}(Xlnb_i)g_{F_i}(V,W)$
$+ \lambda_1\frac{Xb_r}{b_r}\pi(W)V - \lambda_1 g(W,\nabla_V P)X - \lambda_2 g(V,W)\frac{Xb_r}{b_r}P - \lambda_1\lambda_2 g(V,W)\pi(P)X$
$+ \lambda_1^2\pi(W)\pi(V)X$ *if* $i \neq j$;
$(10)\overline{R}(U,V)W = -g(V,W)\frac{g_B(grad_B b_i, grad_B b_k)}{b_i b_k}U - \lambda_1 g(W,\nabla_V P)U - \lambda_2 g(V,W)\nabla_U P -$
$\lambda_1\lambda_2\pi(P)g(V,W)U + \lambda_2^2 g(V,W)\pi(U)P + \lambda_1^2\pi(W)[\pi(V)U - \pi(U)V]$ *if* $i = j \neq k$;
$(11)\overline{R}(U,V)W = g(U,W)grad_B(Vlnb_i) - g(V,W)grad_B(Ulnb_i) + R^{F_i}(U,V)W$
$-\frac{|grad_B b_i|^2_B}{b_i^2}[g(V,W)U - g(U,W)V] + \lambda_1\lambda_2\pi(P)[g(U,W)V - g(V,W)U]$
if $i = j = k \neq r$;
$(12)\overline{R}(U,V)W = g(U,W)grad_B(Vlnb_i) - g(V,W)grad_B(Ulnb_i) + R^{F_i}(U,V)W$
$-\frac{|grad_B f|^2_B}{f^2}[g(V,W)U - g(U,W)V] + \lambda_1[g(W,\nabla_U P)V - g(W,\nabla_V P)U]$
$+ \lambda_2[g(U,W)\nabla_V P - g(V,W)\nabla_U P)] + \lambda_1\lambda_2\pi(P)[g(U,W)V - g(V,W)U]$
$+ \lambda_2^2[g(V,W)\pi(U) - g(U,W)\pi(V)]P + \lambda_1^2\pi(W)[\pi(V)U - \pi(U)V]$ *if* $i = j = k = r$.
where δ_i^r *denotes the Kronecker symbol.*

By Propositions 4.3 and 4.4 and the definition of the Ricci curvature tensor, we have the following two Propositions:

Proposition 4.5. *Let* $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$ *be a multiply twisted product,* $\dim M = \overline{n}$, $\dim B = n$, $\dim F_i = l_i$. *If* $X, Y, Z \in \Gamma(TB)$, $V \in \Gamma(TF_i), W \in \Gamma(TF_j)$ *and* $P \in \Gamma(TB)$, *then:*
$(1)\overline{Ric}(X,Y) = \overline{Ric}^B(X,Y) + \sum_{i=1}^{m} l_i\Big[\frac{H_B^{b_i}(X,Y)}{b_i} + \lambda_2\frac{Pb_i}{b_i}g(X,Y) + \lambda_1\lambda_2\pi(P)g(X,Y)$
$+ \lambda_1 g(Y,\nabla_X P) - \lambda_1^2\pi(X)\pi(Y)\Big]$;
$(2)\overline{Ric}(X,V) = \overline{Ric}(V,X) = (l_i - 1)[VX(lnb_i)]$;
$(3)\overline{Ric}(V,W) = 0$ *if* $i \neq j$;
$(3)\overline{Ric}(V,W) = \overline{Ric}^{F_i}(V,W) + \Big\{\frac{\Delta_B b_i}{b_i} + (l_i - 1)\frac{|grad_B b_i|^2_B}{b_i^2} + \sum_{s\neq i} l_s\frac{g_B(grad_B b_i, grad_B b_s)}{b_i b_s}$
$+[(\overline{n}-1)\lambda_1\lambda_2 - \lambda_2^2]\pi(P) + \lambda_2 div_B P + \lambda_2\sum_{s\neq i} l_s\frac{Pb_s}{b_s} + [(\overline{n}-1)\lambda_1 + (l_i-1)\lambda_2]\frac{Pb_i}{b_i}\Big\}g(V,W)$
if $i = j$.
where $div_B P = \sum_{k=1}^{n}\varepsilon_k\langle\nabla_{E_k}P, E_k\rangle$, *and* $E_k, 1 \leq k \leq n$ *is an orthonormal base of* B *with* $\varepsilon_k = g(E_k, E_k)$.

As a Corollary of Proposition 4.5, we have:

Corollary 4.6. *Let* $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$ *be a multiply twisted product, and* $\dim F_i = l_i > 1$, $P \in \Gamma(TB)$, *then* $(M, \overline{\nabla})$ *is mixed Ricci-flat if and only if* M *can be*

expressed as a multiply warped product. In particular, if $(M, \overline{\nabla})$ *is Einstein, then* M *can be expressed as a multiply warped product.*

Proof. By Proposition 4.5.(2) and (3), similar to the proof of Theorem 1 in [5], we get this Corollary. □

Proposition 4.7. *Let* $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$ *be a multiply twisted product,* $\dim M = \overline{n}$, $\dim B = n$, $\dim F_i = l_i$. *If* $X, Y, Z \in \Gamma(TB)$, $V \in \Gamma(TF_i), W \in \Gamma(TF_j)$ *and* $P \in \Gamma(TF_r)$ *for a fixed* r, *then:*

$(1)\overline{Ric}(X,Y) = \overline{Ric}^B(X,Y) + \sum_{i=1}^{m} l_i \frac{H_B^{b_i}(X,Y)}{b_i} + [(\overline{n}-1)\lambda_1\lambda_2 - \lambda_2^2]\pi(P)g(X,Y)$
$+ \lambda_2 g(X,Y) div_{F_r} P;$

$(2)\overline{Ric}(X,V) = (l_i - 1)[VX(lnb_i)] + [(\overline{n}-1)\lambda_1 - \lambda_2]\pi(V)\frac{Xb_r}{b_r};$

$(3)\overline{Ric}(V,X) = (l_i - 1)[VX(lnb_i)] + [\lambda_2 - (\overline{n}-1)\lambda_1]\pi(V)\frac{Xb_r}{b_r};$

$(4)\overline{Ric}(V,W) = 0$ *if* $i \neq j$;

$(4)\overline{Ric}(V,W) = Ric^{F_i}(V,W) + g(V,W)\Big\{ \frac{\Delta_B b_i}{b_i} + (l_i - 1)\frac{|grad_B b_i|_B^2}{b_i^2} + \sum_{s \neq i} l_s \frac{g_B(grad_B b_i, grad_B b_s)}{b_i b_s}$
$+ [(\overline{n}-1)\lambda_1\lambda_2 - \lambda_2^2]\pi(P)\Big\} + [(\overline{n}-1)\lambda_1 - \lambda_2]g(W, \nabla_V P) + [\lambda_2^2 + (1-\overline{n})\lambda_1^2]\pi(V)\pi(W)$
$+ \lambda_2 g(V,W) div_{F_r} P$ *if* $i = j$.

As a Corollary of Proposition 4.7, we have:

Corollary 4.8. *Let* $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$ *be a multiply twisted product, and* $\dim F_i = l_i > 1$, $P \in \Gamma(TF_r)$, *then* $(M, \overline{\nabla})$ *is mixed Ricci-flat if and only if one of the following two conditions is satisfied:*

(1) $\lambda_2 = (\overline{n}-1)\lambda_1$, *and* M *can be expressed as a multiply warped product;*

(2) $\lambda_2 \neq (\overline{n}-1)\lambda_1$, M *can be expressed as a multiply warped product and* b_r *is only dependent on* F_r;

In particular, if $(M, \overline{\nabla})$ *is Einstein, then* M *can be expressed as a multiply warped product.*

Proof. By Proposition 4.7.(2) and (3), we have that $(M, \overline{\nabla})$ is mixed Ricci-flat if and only if $VX(lnb_i) = 0$, $[(\overline{n}-1)\lambda_1 - \lambda_2]\pi(V)\frac{Xb_r}{b_r} = 0$. Similar to the proof of Corollary 4.6, we get that

$\lambda_2 = (\overline{n}-1)\lambda_1$, and M can be expressed as a multiply warped product.

$\lambda_2 \neq (\overline{n}-1)\lambda_1$, M can be expressed as a multiply warped product. When $i \neq r$, $\pi(V) = 0$. When $i = r$, by $\pi(V)\frac{Xb_r}{b_r} = 0$, then b_r depends only on F_r. □

By Proposition 4.5 and the definition of the scalar curvature, we have the following:

Proposition 4.9. *Let $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$ be a multiply twisted product, $\dim M = \overline{n}$, $\dim B = n$, $\dim F_i = l_i$. If $P \in \Gamma(TB)$, then the scalar curvature $\overline{S}$ has the following expression:*

$$\begin{aligned}\overline{S} = \overline{S}^B &+ 2\sum_{i=1}^{m} l_i \frac{\Delta_B b_i}{b_i} + \sum_{i=1}^{m} \frac{S^{F_i}}{b_i^2} + \sum_{i=1}^{m} l_i(l_i-1)\frac{|grad_B b_i|_B^2}{b_i^2} + \sum_{i=1}^{m}\sum_{s\neq i} l_i l_s \frac{g_B(grad_B b_i, grad_B b_s)}{b_i b_s} \\ &+ \sum_{i=1}^{m} l_i[(\overline{n}-1)\lambda_1 + (n+l_i-1)\lambda_2)]\frac{Pb_i}{b_i} + \lambda_2 \sum_{i=1}^{m}\sum_{s\neq i} l_i l_s \frac{Pb_s}{b_s} + \sum_{i=1}^{m} l_i[(\overline{n}+n-1)\lambda_1\lambda_2 \\ &- (\lambda_1^2+\lambda_2^2)]\pi(P) + (\lambda_1+\lambda_2)\sum_{i=1}^{m} l_i div_B P.\end{aligned}$$

By Proposition 4.7 and the definition of the scalar curvature, we have the following:

Proposition 4.10. *Let $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$ be a multiply twisted product, $\dim M = \overline{n}$, $\dim B = n$, $\dim F_i = l_i$. If $P \in \Gamma(TF_r)$, then the scalar curvature $\overline{S}$ has the following expression:*

$$\begin{aligned}\overline{S} = \overline{S}^B &+ 2\sum_{i=1}^{m} l_i \frac{\Delta_B b_i}{b_i} + \sum_{i=1}^{m} \frac{S^{F_i}}{b_i^2} + \sum_{i=1}^{m} l_i(l_i-1)\frac{|grad_B b_i|_B^2}{b_i^2} + \sum_{i=1}^{m}\sum_{s\neq i} l_i l_s \frac{g_B(grad_B b_i, grad_B b_s)}{b_i b_s} \\ &+ [\overline{n}(\overline{n}-1)\lambda_1\lambda_2 + (1-\overline{n})(\lambda_1^2+\lambda_2^2)]\pi(P) + (\overline{n}-1)(\lambda_1+\lambda_2) div_{F_r} P.\end{aligned}$$

Remark 11. (1) *It is easy to see that Propositions* $3.1 - 3.8$ *are Corollaries of Propositions* $4.1, 4.2, 4.3, 4.4, 4.5, 4.7, 4.9, 4.10$, *respectively.*

(2) *When* $\lambda_1 = \lambda_2 = 1$, *we get Propositions* $1, 2, 4, 5, 7, 9, 12, 13$ *in* [12], *by Propositions* $4.1, 4.2, 4.3, 4.4, 4.5, 4.7, 4.9, 4.10$, *respectively.*

4.2 Special Multiply Warped Product with a Quarter-symmetric Connection

Let $M = I \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$ be a multiply warped product with the metric tensor $g = -dt^2 \oplus b_1^2 g_{F_1} \oplus b_2^2 g_{F_2} \cdots \oplus b_m^2 g_{F_m}$ and I is an open interval in $\mathbb{R}$, and $b_i \in C^\infty(I)$, $\dim M = \overline{n}$, $\dim I = 1$, $\dim F_i = l_i$.

Similar to the proof method of Theorem 3.11, we have:

Theorem 4.11. *Let $M = I \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$ be a multiply warped product with the metric tensor $g = -dt^2 \oplus b_1^2 g_{F_1} \oplus b_2^2 g_{F_2} \cdots \oplus b_m^2 g_{F_m}$, $P = \frac{\partial}{\partial t}$. Then $(M, \overline{\nabla})$ is Einstein with the Einstein constant α if and only if the following conditions are satisfied*

(1) *(F_i, ∇^{F_i}) is Einstein with the Einstein constant α_i, $i \in \{1, \ldots, m\}$;*

(2) $\sum_{i=1}^{m} l_i\Big(\lambda_2\frac{b_i'}{b_i} - \frac{b_i''}{b_i} + \lambda_1^2 - \lambda_1\lambda_2\Big) = \alpha$;

(3) $\alpha_i - b_i b_i'' + (1-l_i)(b_i')^2 + (\lambda_2 b_i^2 - b_i b_i')\sum\limits_{s\neq i} l_s \frac{b_s'}{b_s} + [\lambda_2^2 + (1-\overline{n})\lambda_1\lambda_2]b_i^2 + [(\overline{n}-1)\lambda_1 + (l_i-1)\lambda_2]b_i b_i' = \alpha b_i^2$.

Theorem 4.12. *Let $M = I \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$ be a multiply warped product with the metric tensor $g = -dt^2 \oplus b_1^2 g_{F_1} \oplus b_2^2 g_{F_2} \cdots \oplus b_m^2 g_{F_m}$, $P \in \Gamma(TF_r)$ with $g_{F_r}(P,P) = 1$ and $\overline{n} > 2$. Then $(M, \overline{\nabla})$ is Einstein with the Einstein constant α if and only if the following conditions are satisfied for any $i \in \{1, \ldots, m\}$:*

(1) *$(F_i, \nabla^{F_i})(i \neq r)$ is Einstein with the Einstein constant $\alpha_i, i \in \{1, \ldots, m\}$;*

(2) *b_r is a constant and $\sum\limits_{i=1}^{m} l_i \frac{b_i''}{b_i} = \mu_0$; $div_{F_r} P = \mu_1$; $\mu_0 - \lambda_2\mu_1 + \alpha = [(\overline{n}-1)\lambda_1\lambda_2 - \lambda_2^2]b_r^2$, where μ_0, μ_1 are constants;*

(3) *$Ric^{F_r}(V,W) + \overline{\alpha} g_{F_r}(V,W) = [(\overline{n}-1)\lambda_1^2 - \lambda_2^2]\pi(V)\pi(W) - [(\overline{n}-1)\lambda_1 - \lambda_2]g(W, \nabla_V P)$, for $V, W \in \Gamma(TF_r)$;*

(4) *$\alpha_i - b_i b_i'' + [(\overline{n}-1)\lambda_1\lambda_2 - \lambda_2^2]b_i^2 b_r^2 - b_i b_i' \sum\limits_{s\neq i} l_s \frac{b_s'}{b_s} - (l_i - 1)(b_i')^2 = (\alpha - \lambda_2\mu_1)b_i^2$.*

Proof. By Proposition 4.7(2) and $g_{F_r}(P,P) = 1$, we have that b_r is a constant. By Proposition 4.7(1), we have

$$\overline{Ric}\Big(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\Big) = \sum_{i=1}^{m} l_i \frac{b_i''}{b_i} + [\lambda_2^2 + (1-\overline{n})\lambda_1\lambda_2]b_r^2 - \lambda_2 div_{F_r} P = -\alpha.$$

By variables separation, we have

$$\sum_{i=1}^{m} l_i \frac{b_i''}{b_i} = \mu_0;\ div_{F_r} P = \mu_1;\ \mu_0 - \lambda_2\mu_1 + \alpha = [(\overline{n}-1)\lambda_1\lambda_2 - \lambda_2^2]b_r^2,$$

then we get (2). By Proposition 4.7(3), we have

$$\begin{aligned}
\overline{Ric}(V,W) &= Ric^{F_i}(V,W) + b_i^2 g_{F_i}(V,W)\Big\{ -\frac{b_i''}{b_i} + (l_i-1)\frac{-(b_i')^2}{b_i^2} + \sum_{s\neq i} l_s \frac{-b_i' b_s'}{b_i b_s} \\
&\quad + [(\overline{n}-1)\lambda_1\lambda_2 - \lambda_2^2]\pi(P)\Big\} + [(\overline{n}-1)\lambda_1 - \lambda_2]g(W, \nabla_V P) \\
&\quad + [\lambda_2^2 + (1-\overline{n})\lambda_1^2]\pi(V)\pi(W) + \lambda_2 g(V,W) div_{F_r} P.
\end{aligned}$$

When $i \neq r$, then $\nabla_V P = \pi(V) = 0$, so

$$\begin{aligned}
\overline{Ric}(V,W) &= Ric^{F_i}(V,W) + b_i^2 g_{F_i}(V,W)\Big\{ -\frac{b_i''}{b_i} + (l_i-1)\frac{-(b_i')^2}{b_i^2} + \sum_{s\neq i} l_s \frac{-b_i' b_s'}{b_i b_s} \\
&\quad + [(\overline{n}-1)\lambda_1\lambda_2 - \lambda_2^2]b_r^2\Big\} + \lambda_2\mu_1 b_i^2 g_{F_i}(V,W) = \alpha b_i^2 g_{F_i}(V,W).
\end{aligned}$$

By variables separation, we have that $(F_i, \nabla^{F_i})(i \neq r)$ is Einstein with the Einstein constant α_i and

$$\alpha_i - b_i b_i'' + [(\overline{n}-1)\lambda_1\lambda_2 - \lambda_2^2]b_i^2 b_r^2 - b_i b_i' \sum_{s\neq i} l_s \frac{b_s'}{b_s} - (l_i - 1)(b_i')^2 = (\alpha - \lambda_2\mu_1)b_i^2.$$

Then we get (1) and (4).
When $i = r$ and b_r is a constant, then

$$\begin{aligned} &Ric^{F_r}(V,W) + b_r^2\big\{[(\overline{n}-1)\lambda_1\lambda_2 - \lambda_2^2]b_r^2 + \lambda_2\mu_1 - \alpha\big\}g_{F_r}(V,W) \\ &\quad = [(\overline{n}-1)\lambda_1^2 - \lambda_2^2]\pi(V)\pi(W) - [(\overline{n}-1)\lambda_1 - \lambda_2]g(W, \nabla_V P), \end{aligned}$$

let $\overline{\alpha} = b_r^2\big\{[(\overline{n}-1)\lambda_1\lambda_2 - \lambda_2^2]b_r^2 + \lambda_2\mu_1 - \alpha\big\}$, we get (3). □

Theorem 4.13. *Let $M = I \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$ be a multiply warped product and $P = \frac{\partial}{\partial t}$. If $(M, \overline{\nabla})$ has constant scalar curvature $\overline{S}$, then each (F_i, ∇^{F_i}) has constant scalar curvature S^{F_i}.*

Proof. By Proposition 4.9, we have

$$\begin{aligned} \overline{S} \;=\; & -2\sum_{i=1}^m l_i \frac{b_i''}{b_i} + \sum_{i=1}^m \frac{S^{F_i}}{b_i^2} + \sum_{i=1}^m l_i(l_i-1)\frac{-(b_i')^2}{b_i^2} + \sum_{i=1}^m \sum_{s\neq i} l_i l_s \frac{-b_i' b_s'}{b_i b_s} \\ & + \sum_{i=1}^m l_i[(\overline{n}-1)\lambda_1 + l_i\lambda_2)]\frac{b_i'}{b_i} + \lambda_2 \sum_{i=1}^m \sum_{s\neq i} l_i l_s \frac{b_s'}{b_s} \\ & - \sum_{i=1}^m l_i[\overline{n}\lambda_1\lambda_2 - (\lambda_1^2 + \lambda_2^2)]. \end{aligned} \tag{38}$$

Note that each S^{F_i} is function defined on F_i, using variables separation we complete this proof. □

Theorem 4.14. *Let $M = I \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$ be a multiply warped product and $P \in \Gamma(TF_r)$. If $(M, \overline{\nabla})$ has constant scalar curvature $\overline{S}$, then*
(1) each $(F_i, \nabla^{F_i})(i \neq r)$ has constant scalar curvature S^{F_i};
(2) If $\lambda_1 + \lambda_2 \neq 0$ and $\lambda_1^2 + \lambda_2^2 - \overline{n}\lambda_1\lambda_2 = 0$, and $div_{F_r} P$ is a constant, then S^{F_r} is a constant;
(3) If $\lambda_1 = -\lambda_2 \neq 0$ and $g_{F_r}(P,P)$ is a constant, then S^{F_r} is a constant;
(4) If $\lambda_1 + \lambda_2 \neq 0$ and $\lambda_1^2 + \lambda_2^2 - \overline{n}\lambda_1\lambda_2 \neq 0$, and $div_{F_r} P, g_{F_r}(P,P)$ are constants, then S^{F_r} is a constant.

Proof. By Proposition 4.10, we have

$$\begin{aligned}\overline{S} &= -2\sum_{i=1}^{m} l_i\frac{b_i''}{b_i}+\sum_{i=1}^{m}\frac{S^{F_i}}{b_i^2}+\sum_{i=1}^{m} l_i(l_i-1)\frac{-(b_i')^2}{b_i^2}+\sum_{i=1}^{m}\sum_{s\neq i} l_i l_s\frac{-b_i'b_s'}{b_ib_s} \\ &\quad +[\overline{n}(\overline{n}-1)\lambda_1\lambda_2+(1-\overline{n})(\lambda_1^2+\lambda_2^2)]b_r^2 g_{F_r}(P,P)+(\overline{n}-1)(\lambda_1+\lambda_2)div_{F_r}P \end{aligned} \tag{39}$$

then similar to the proof of Proposition 3.18, we complete this Proposition. □

Remark 12. *When $\lambda_1=\lambda_2=1$, we get Theorems* 15, 16, *Propositions* 18, 19 *in* [12] *by Theorems* $4.11-4.14$, *respectively.*

4.3 Generalized Kasner Space-times with a Quarter-symmetric Connection

In this section, we consider the Einstein and scalar curvature of generalized Kasner space-times with a quarter-symmetric connection. We recall the definition of generalized Kasner space-times in [5].

Definition 4.15. *A generalized Kasner space-time (M,g) is a Lorentzian multiply warped product of the form $M=I\times_{\phi^{p_1}}F_1\times_{\phi^{p_2}}F_2\cdots\times_{\phi^{p_m}}F_m$ with the metric tensor $g=-dt^2\oplus\phi^{2p_1}g_{F_1}\oplus\phi^{2p_2}g_{F_2}\cdots\oplus\phi^{2p_m}g_{F_m}$, where $\phi:I\to(0,\infty)$ is smooth and $p_i\in\mathbb{R}$, for any $i\in\{1,\ldots,m\}$ and also $I=(t_1,t_2)$.*

We introduce the following parameters $\zeta=\sum_{i=1}^{m} l_ip_i$ and $\eta=\sum_{i=1}^{m} l_ip_i^2$ for generalized Kasner space-times. By Theorem 4.11 and direct computations, we get the following:

Proposition 4.16. *Let $M=I\times_{\phi^{p_1}}F_1\times_{\phi^{p_2}}F_2\cdots\times_{\phi^{p_m}}F_m$ be a generalized Kasner space-time and $P=\frac{\partial}{\partial t}$. Then $(M,\overline{\nabla})$ is Einstein with the Einstein constant α if and only if the following conditions are satisfied for any $i\in\{1,\ldots,m\}$:*
(1) *(F_i,∇^{F_i}) is Einstein with the Einstein constant $\alpha_i, i\in\{1,\ldots,m\}$;*
(2) $\zeta\Big(\lambda_2\frac{\phi'}{\phi}-\frac{\phi''}{\phi}\Big)-(\eta-\zeta)\frac{(\phi')^2}{\phi^2}+(\lambda_1^2-\lambda_1\lambda_2)(\overline{n}-1)=\alpha$;
(3) $\frac{\alpha_i}{\phi^{2p_i}}-p_i\frac{\phi''}{\phi}-(\zeta-1)p_i\frac{(\phi')^2}{\phi^2}+\{\lambda_2\zeta+[(\overline{n}-1)\lambda_1-\lambda_2]p_i\}\frac{\phi'}{\phi}=\alpha-\lambda_2^2+(\overline{n}-1)\lambda_1\lambda_2$.

By the equation (38), we obtain the following:

Proposition 4.17. *Let $M=I\times_{\phi^{p_1}}F_1\times_{\phi^{p_2}}F_2\cdots\times_{\phi^{p_m}}F_m$ be a generalized Kasner space-time and $P=\frac{\partial}{\partial t}$. Then $(M,\overline{\nabla})$ has constant scalar curvature $\overline{S}$ if and only if*

each (F_i, ∇^{F_i}) *has constant scalar curvature* S^{F_i} *and*

$$\overline{S} = \sum_{i=1}^{m} \frac{S^{F_i}}{\phi^{2p_i}} - 2\zeta\frac{\phi''}{\phi} - (\eta+\zeta^2-2\zeta)\frac{(\phi')^2}{\phi^2} + (\lambda_1+\lambda_2)\zeta(\overline{n}-1)\frac{\phi'}{\phi} + (\overline{n}-1)(\lambda_1^2+\lambda_2^2-\overline{n}\lambda_1\lambda_2). \tag{40}$$

Next, we first give a classification of four-dimensional generalized Kasner space-times with a quarter-symmetric connection and then consider Ricci tensors and scalar curvatures of them.

Definition 4.18. *Let* $M = I \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$ *be a multiply warped product with the metric tensor* $g = -dt^2 \oplus b_1^2 g_{F_1} \oplus b_2^2 g_{F_2} \cdots \oplus b_m^2 g_{F_m}$. *Then:*
(1) (M, g) *is said to be of type* (I) *if* $m = 1$ *and* $\dim F = 3$;
(2) (M, g) *is said to be of type* (II) *if* $m = 2$ *and* $\dim F_1 = 1$ *and* $\dim F_2 = 2$;
(3) (M, g) *is said to be of type* (III) *if* $m = 3$ *and* $\dim F_1 = 1,\ \dim F_2 = 1,\ \dim F_3 = 1$.

4.3.1 Classification of Einstein Type (I) generalized Kasner space-times with a quarter-symmetric connection

By Theorem 3.16, we have given a classification of Einstein Type (I) generalized Kasner space-times with a quarter-symmetric connection.

4.3.2 Type (I) generalized Kasner space-times with a quarter-symmetric connection with constant scalar curvature

By Theorem 3.19, we have given a classification of Type (I) generalized Kasner space-times with a quarter-symmetric connection with constant scalar curvature.

4.3.3 Classification of Einstein Type (II) generalized Kasner space-times with a quarter-symmetric connection

Let $M = I \times_{\phi^{p_1}} F_1 \times_{\phi^{p_2}} F_2$ be an Einstein Type (II) generalized Kasner space-times and $P = \frac{\partial}{\partial t}$. Then $\alpha_1 = 0$ because of $\dim F_1 = 1$. $\zeta = p_1 + 2p_2,\ \eta = p_1^2 + 2p_2^2$. By Proposition 4.16, we have

$$\zeta\Big(\lambda_2\frac{\phi'}{\phi} - \frac{\phi''}{\phi}\Big) - (\eta-\zeta)\frac{(\phi')^2}{\phi^2} + 3(\lambda_1^2 - \lambda_1\lambda_2) = \alpha. \tag{41a}$$

$$-p_1\frac{\phi''}{\phi}-(\zeta-1)p_1\frac{(\phi')^2}{\phi^2}+[\lambda_2\zeta+(3\lambda_1-\lambda_2)p_1]\frac{\phi'}{\phi}=\alpha-\lambda_2^2+3\lambda_1\lambda_2. \tag{41b}$$

$$\frac{\alpha_2}{\phi^{2p_2}}-p_2\frac{\phi''}{\phi}-(\zeta-1)p_2\frac{(\phi')^2}{\phi^2}+[\lambda_2\zeta+(3\lambda_1-\lambda_2)p_2]\frac{\phi'}{\phi}=\alpha-\lambda_2^2+3\lambda_1\lambda_2. \tag{41c}$$

where α_2 is a constant. Consider the following two cases:

Case $\underline{(i)\ (\zeta=0).}$ In this case, $p_2=-\frac{1}{2}p_1,\ \eta=\frac{3}{2}p_1^2$. Then by equations $(41a)-(41c)$, we have:

$$-\eta\frac{(\phi')^2}{\phi^2}+3(\lambda_1^2-\lambda_1\lambda_2)=\alpha. \tag{42a}$$

$$p_1\Big[-\frac{\phi''}{\phi}+\frac{(\phi')^2}{\phi^2}+(3\lambda_1-\lambda_2)\frac{\phi'}{\phi}\Big]=\alpha-\lambda_2^2+3\lambda_1\lambda_2. \tag{42b}$$

$$\frac{\alpha_2}{\phi^{-p_1}}-\frac{1}{2}p_1\Big[-\frac{\phi''}{\phi}+\frac{(\phi')^2}{\phi^2}+(3\lambda_1-\lambda_2)\frac{\phi'}{\phi}\Big]=\alpha-\lambda_2^2+3\lambda_1\lambda_2. \tag{42c}$$

$\underline{\eta=0.}$ We have $p_i=0$, by the equation $(42a)$, we get $\alpha=3\lambda_1^2-3\lambda_1\lambda_2$; by the equation $(42b)$, we get $\alpha=\lambda_2^2-3\lambda_1\lambda_2$; then we have $\lambda_2^2=3\lambda_1^2$ and by the equation $(42c)$, we get $\alpha_2=0$. So we have $\underline{\lambda_2^2=3\lambda_1^2,\ p_i=0,\ \alpha=3\lambda_1^2-3\lambda_1\lambda_2=\lambda_2^2-3\lambda_1\lambda_2,\ \alpha_1=\alpha_2=0.}$

$\underline{\eta\neq0.}$ We have $p_i\neq0$.

1) $\underline{\alpha_2=0.}$ By equations $(42b),(42c)$, we get $\alpha=\lambda_2^2-3\lambda_1\lambda_2$ and

$$-\frac{\phi''}{\phi}+\frac{(\phi')^2}{\phi^2}+(3\lambda_1-\lambda_2)\frac{\phi'}{\phi}=0; \tag{43a}$$

$$\frac{(\phi')^2}{\phi^2}=\frac{3\lambda_1^2-\lambda_2^2}{\eta}. \tag{43b}$$

$1'.\ 3\lambda_1^2-\lambda_2^2<0$, we have no solution;

$2'.\ 3\lambda_1^2-\lambda_2^2=0$, we have $\phi=c$, then by the equation $(42b)$, we get $\alpha=\lambda_2^2-3\lambda_1\lambda_2$.

So we have $\underline{\lambda_2^2=3\lambda_1^2,\ p_1\neq0,\ p_2\neq0\ \alpha=\lambda_2^2-3\lambda_1\lambda_2,\ \alpha_1=\alpha_2=0,\ \phi=c.}$

$3'.\ 3\lambda_1^2-\lambda_2^2>0$, we have $\phi=c_0e^{\pm\sqrt{\frac{3\lambda_1^2-\lambda_2^2}{\eta}}t}$, by the equation $(43a)$, we get $\lambda_2=3\lambda_1$, considering that $3\lambda_1^2-\lambda_2^2>0$, we have $\lambda_1^2<0$, which is a contradiction.

2) $\underline{\alpha_2\neq0.}$ By equations $(42b),(42c)$, we get $\frac{\alpha_2}{\phi^{-p_1}}=\frac{3}{2}(\alpha-\lambda_2^2+3\lambda_1\lambda_2)$, so $\phi=c$; then by the equation $(42b)$, we get $\alpha-\lambda_2^2+3\lambda_1\lambda_2=0$, then $\alpha_2=0$, this is a contradiction.

Case $\underline{(ii)\ (\zeta\neq0).}$ Then $\eta\neq0$. Putting $\phi=\psi^{\frac{\zeta}{\eta}}$, then $\psi''-\lambda_2\psi'+(\alpha+3\lambda_1\lambda_2-3\lambda_1^2)\frac{\eta}{\zeta^2}\psi=0$. Hence:

(1) $\alpha<\frac{\lambda_2^2\zeta^2}{4\eta}+3\lambda_1^2-3\lambda_1\lambda_2,\ \psi=c_1e^{\frac{\lambda_2+\sqrt{\lambda_2^2-4(\alpha+3\lambda_1\lambda_2-3\lambda_1^2)\frac{\eta}{\zeta^2}}}{2}t}+c_2e^{\frac{\lambda_2-\sqrt{\lambda_2^2-4(\alpha+3\lambda_1\lambda_2-3\lambda_1^2)\frac{\eta}{\zeta^2}}}{2}t}$;

(2) $\alpha=\frac{\lambda_2^2\zeta^2}{4\eta}+3\lambda_1^2-3\lambda_1\lambda_2,\ \psi=c_1e^{\frac{\lambda_2}{2}t}+c_2te^{\frac{\lambda_2}{2}t}$;

(3) $\alpha > \frac{\lambda_2^2\zeta^2}{4\eta} + 3\lambda_1^2 - 3\lambda_1\lambda_2$,

$\psi = c_1 e^{\frac{\lambda_2}{2}t} cos(\frac{\sqrt{4(\alpha+3\lambda_1\lambda_2-3\lambda_1^2)\frac{\eta}{\zeta^2}-\lambda_2^2}}{2}t) + c_2 e^{\frac{\lambda_2}{2}t} sin(\frac{\sqrt{4(\alpha+3\lambda_1\lambda_2-3\lambda_1^2)\frac{\eta}{\zeta^2}-\lambda_2^2}}{2}t)$.

We make equations $(41a)-(41c)$ into

$$\frac{\zeta^2}{\eta}\frac{\lambda_2\psi' - \psi''}{\psi} = \alpha + 3\lambda_1\lambda_2 - 3\lambda_1^2; \tag{44a}$$

$$-\frac{p_1}{\zeta}\frac{(\phi^\zeta)''}{\phi^\zeta} + \frac{\lambda_2\zeta + (3\lambda_1-\lambda_2)p_1}{\zeta}\frac{(\phi^\zeta)'}{\phi^\zeta} = \alpha - \lambda_2^2 + 3\lambda_1\lambda_2; \tag{44b}$$

$$\frac{\alpha_2}{\phi^{2p_2}} - \frac{p_2}{\zeta}\frac{(\phi^\zeta)''}{\phi^\zeta} + \frac{\lambda_2\zeta + (3\lambda_1-\lambda_2)p_2}{\zeta}\frac{(\phi^\zeta)'}{\phi^\zeta} = \alpha - \lambda_2^2 + 3\lambda_1\lambda_2. \tag{44c}$$

When $p_1 = p_2$, type (II) spaces turn into type (I) spaces, so we assume $p_1 \neq p_2$. By $(44b)\times p_2 - (44c)\times p_1$, we get

$$\psi' = \frac{p_1\alpha_2}{\lambda_2(p_2-p_1)}\frac{\eta}{\zeta^2}\psi^{1-\frac{2p_2\zeta}{\eta}} + \frac{\alpha-\lambda_2^2+3\lambda_1\lambda_2}{\lambda_2}\frac{\eta}{\zeta^2}\psi. \tag{45}$$

$\alpha < \frac{\lambda_2^2\zeta^2}{4\eta} + 3\lambda_1^2 - 3\lambda_1\lambda_2$, $\psi = c_1e^{at} + c_2e^{bt}$, where $a = \frac{\lambda_2+\sqrt{\lambda_2^2-4(\alpha+3\lambda_1\lambda_2-3\lambda_1^2)\frac{\eta}{\zeta^2}}}{2}$, $b = \frac{\lambda_2-\sqrt{\lambda_2^2-4(\alpha+3\lambda_1\lambda_2-3\lambda_1^2)\frac{\eta}{\zeta^2}}}{2}$. By the equation (45), we get

$$ac_1e^{at} + bc_2e^{bt} = \frac{p_1\alpha_2}{\lambda_2(p_2-p_1)}\frac{\eta}{\zeta^2}\big(c_1e^{at}+c_2e^{bt}\big)^{1-\frac{2p_2\zeta}{\eta}} + \frac{\alpha-\lambda_2^2+3\lambda_1\lambda_2}{\lambda_2}\frac{\eta}{\zeta^2}\big(c_1e^{at}+c_2e^{bt}\big). \tag{46}$$

1) $\underline{c_1 = 0.}$ We have

$$\Big[b - \frac{\alpha-\lambda_2^2+3\lambda_1\lambda_2}{\lambda_2}\frac{\eta}{\zeta^2}\Big]c_2e^{bt} = \frac{p_1\alpha_2}{\lambda_2(p_2-p_1)}\frac{\eta}{\zeta^2}\big(c_2e^{bt}\big)^{1-\frac{2p_2\zeta}{\eta}} \tag{47}$$

$1'$. $\underline{b\neq 0,\ p_1\alpha_2 \neq 0.}$ By the equation (47), we get $p_2 = 0$, so $\zeta = p_1$, $\eta = p_1^2$ and $\frac{\zeta^2}{\eta} = 1$. Then $b = \frac{\lambda_2-\sqrt{\lambda_2^2-4(\alpha+3\lambda_1\lambda_2-3\lambda_1^2)}}{2}$, so

$$b^2 - \lambda_2 b = 3\lambda_1^2 - 3\lambda_2\lambda_2 - \alpha. \tag{48}$$

On the other hand, $\phi^\zeta = \psi = c_2e^{bt}$, so by equations $(44b), (44c)$, we get

$$-b^2 + 3\lambda_1 b = \alpha - \lambda_2^2 + 3\lambda_1\lambda_2; \tag{49a}$$

$$\alpha_2 + \lambda_2 b = \alpha - \lambda_2^2 + 3\lambda_1\lambda_2; \tag{49b}$$

$(48)+(49a)$, we get $(3\lambda_1-\lambda_2)b=3\lambda_1^2-\lambda_2^2$, if $\lambda_2=3\lambda_1$, then $0=-6\lambda_1^2$, this is a contradiction with $\lambda_1\neq 0$, so $\lambda_2\neq 3\lambda_1$, $b=\frac{3\lambda_1^2-\lambda_2^2}{3\lambda_1-\lambda_2}$. Then by the equation (48), we have $\alpha=\frac{18\lambda_1^4-36\lambda_1^3\lambda_2+24\lambda_1^2\lambda_2^2-6\lambda_1\lambda_2^3}{(3\lambda_1-\lambda_2)^2}=\frac{6\lambda_1(\lambda_1-\lambda_2)(3\lambda_1^2-3\lambda_1\lambda_2+\lambda_2^2)}{(3\lambda_1-\lambda_2)^2}$, by the equation $(49b)$, we get $\alpha_2=\frac{18\lambda_1^4-18\lambda_1^3\lambda_2+6\lambda_1\lambda_2^3-2\lambda_2^4}{(3\lambda_1-\lambda_2)^2}=\frac{2(3\lambda_1^2-\lambda_2^2)(3\lambda_1^2-3\lambda_1\lambda_2+\lambda_2^2)}{(3\lambda_1-\lambda_2)^2}$, since $3\lambda_1^2-3\lambda_1\lambda_2+\lambda_2^2\neq 0$, and $\alpha_2\neq 0$, we have $3\lambda_1^2\neq\lambda_2^2$. Considering that $b<\frac{\lambda_2}{2}$, we get $\frac{6\lambda_1^2-3\lambda_1\lambda_2-\lambda_2^2}{3\lambda_1-\lambda_2}<0$.

So we obtain $\underline{3\lambda_1^2\neq\lambda_2^2,\ \frac{6\lambda_1^2-3\lambda_1\lambda_2-\lambda_2^2}{3\lambda_1-\lambda_2}<0,\ p_1\neq 0,\ p_2=0,\ \alpha=\frac{18\lambda_1^4-36\lambda_1^3\lambda_2+24\lambda_1^2\lambda_2^2-6\lambda_1\lambda_2^3}{(3\lambda_1-\lambda_2)^2},\ \alpha_1=0,\ \alpha_2=\frac{18\lambda_1^4-18\lambda_1^3\lambda_2+6\lambda_1\lambda_2^3-2\lambda_2^4}{(3\lambda_1-\lambda_2)^2},\ \phi=c_0e^{\frac{3\lambda_1^2-\lambda_2^2}{(3\lambda_1-\lambda_2)p_1}t}.}$

$\underline{2'.\ b\neq 0,\ p_1\alpha_2=0.}$

1°. $\underline{p_1=0.}$ Then $\zeta=2p_2,\ \eta=2p_2^2,\ \frac{\eta}{\zeta^2}=\frac{1}{2}$. By the equation (47), we get $b=\frac{\alpha-\lambda_2^2+3\lambda_1\lambda_2}{2\lambda_2}$, on the other hand, $b=\frac{\lambda_2-\sqrt{\lambda_2^2-2(\alpha+3\lambda_1\lambda_2-3\lambda_1^2)}}{2}$, so we get

$$\alpha^2+(6\lambda_1\lambda_2-2\lambda_2^2)\alpha+3\lambda_1^2\lambda_2^2-6\lambda_1^3\lambda_2+3\lambda_1^4=0. \tag{50}$$

when $\lambda_1=\lambda_2$, we get $\alpha^2+4\lambda_1^2\alpha=0$, then $\alpha=0$ or $\alpha=-4\lambda_1^2$. If $\alpha=0$, we have $b=\lambda_1=\lambda_2<0$, by the equation $(44c)$, we get $\alpha_2=0$; $\phi=\psi^{\frac{\zeta}{\eta}}=c_0e^{\frac{\lambda_1}{p_2}t}$. If $\alpha=-4\lambda_1^2$, then $\lambda_1=\lambda_2>0,\ b=-\lambda_1$, then by the equation $(44c)$, we get $\lambda_1=0$, which is a contradiction.

When $\lambda_1\neq\lambda_2,\ \Delta=8\lambda_2^2(3\lambda_1^2-\lambda_2^2)$, if $\lambda_2^2>3\lambda_1^2$, we have no solution; if $\lambda_2^2=3\lambda_1^2$, we have $b=0$, which is a contradiction; if $\lambda_2^2<3\lambda_1^2$, we have $\alpha=\lambda_2^2-3\lambda_1\lambda_2+\sqrt{2\lambda_2^2(3\lambda_1^2-\lambda_2^2)}$ or $\alpha=\lambda_2^2-3\lambda_1\lambda_2-\sqrt{2\lambda_2^2(3\lambda_1^2-\lambda_2^2)}$. By the equation $(44c)$, we have $\alpha_2=0$ and

$$-2b^2+(3\lambda_1+\lambda_2)b=\alpha-\lambda_2^2+3\lambda_1\lambda_2. \tag{51}$$

When $\alpha=\lambda_2^2-3\lambda_1\lambda_2+\sqrt{2\lambda_2^2(3\lambda_1^2-\lambda_2^2)}$, we have $b=\frac{\sqrt{2\lambda_2^2(3\lambda_1^2-\lambda_2^2)}}{2\lambda_2}$, by the equation (51), we have $\lambda_1=\lambda_2$, which is a contradiction. When $\alpha=\lambda_2^2-3\lambda_1\lambda_2-\sqrt{2\lambda_2^2(3\lambda_1^2-\lambda_2^2)}$, we have the same contradiction.

So we get $\underline{\lambda_1=\lambda_2<0,\ p_1=0,\ p_2\neq 0,\ \alpha=\alpha_1=\alpha_2=0,\ \phi=c_0e^{\frac{\lambda_1}{p_2}t}.}$

2°. $\underline{\alpha_2=0.}$ Since $p_1=0$ we have discussed, so we assume that $p_1\neq 0$. By the equation (47), we have $b=\frac{\alpha-\lambda_2^2+3\lambda_1\lambda_2}{\lambda_2}\frac{\eta}{\zeta^2},\ \phi^\zeta=ce^{b\frac{\zeta^2}{\eta}t}$, then by the equation$(44b)$, we have $\alpha(\alpha-\lambda_2^2+3\lambda_1\lambda_2)=0$. So when $\lambda_2=3\lambda_1$, we have $\alpha=0$; when $\lambda_2\neq 3\lambda_1$, we have $\alpha=0$ or $\alpha=\lambda_2^2-3\lambda_1\lambda_2\neq 0$.

When $\lambda_2=3\lambda_1,\ \alpha=0$, then $b=0$, which is a contradiction;

when $\lambda_2\neq 3\lambda_1,\ \alpha=\lambda_2^2-3\lambda_1\lambda_2\neq 0$, then $b=0$, which is a contradiction;

when $\lambda_2\neq 3\lambda_1,\ \alpha=0$, then $b=(3\lambda_1-\lambda_2)\frac{\eta}{\zeta^2}$, by the equation $(44a)$, we get $\frac{\eta}{\zeta^2}=\frac{3\lambda_1^2-\lambda_2^2}{(3\lambda_1-\lambda_2)^2}$, then $b=\frac{3\lambda_1^2-\lambda_2^2}{3\lambda_1-\lambda_2}$. Since $b<\frac{\lambda_2}{2}$, we get $\frac{6\lambda_1^2-3\lambda_1\lambda_2-\lambda_2^2}{3\lambda_1-\lambda_2}<0$, by $\alpha<$

$\frac{\lambda_2^2\zeta^2}{4\eta}+3\lambda_1^2-3\lambda_1\lambda_2$, we get $\lambda_2^2<3\lambda_1^2$. $\phi=\psi^{\frac{\zeta}{\eta}}=ce^{\frac{3\lambda_1-\lambda_2}{\zeta}t}$.
So we have $\underline{\lambda_2^2<3\lambda_1^2,\ \frac{6\lambda_1^2-3\lambda_1\lambda_2-\lambda_2^2}{3\lambda_1-\lambda_2}<0,\ p_1\neq 0,\ p_2\neq-\frac{1}{2}p_1,\ \alpha=\alpha_1=\alpha_2=0,\ \phi=ce^{\frac{3\lambda_1-\lambda_2}{\zeta}t}}$.
$3'$. $\underline{b=0.}$ Then $\psi=c_2$, by the equation $(44a)$, we have $\alpha=3\lambda_1^2-3\lambda_1\lambda_2$; by the equation $(44b)$, we have $\alpha=\lambda_2^2-3\lambda_1\lambda_2$; so $\lambda_2^2=3\lambda_1^2$, by the equation $(44c)$, we have $\alpha_2=0$.

So we have $\underline{\lambda_2^2=3\lambda_1^2,\ \zeta\neq 0,\ \eta\neq 0,\ \alpha=\lambda_2^2-3\lambda_1\lambda_2=3\lambda_1^2-3\lambda_1\lambda_2,\ \alpha_1=\alpha_2=0,\ \phi=c.}$
2) $\underline{c_2=0.}$ We have

$$\Big[a-\frac{\alpha-\lambda_2^2+3\lambda_1\lambda_2}{\lambda_2}\frac{\eta}{\zeta^2}\Big]c_1e^{at}=\frac{p_1\alpha_2}{\lambda_2(p_2-p_1)}\frac{\eta}{\zeta^2}\big(c_1e^{at}\big)^{1-\frac{2p_2\zeta}{\eta}}\tag{52}$$

$1'$. $\underline{p_1\alpha_2\neq 0.}$ By the equation (52), we get $p_2=0$, so $\zeta=p_1,\ \eta=p_1^2$ and $\frac{\zeta^2}{\eta}=1$. Then $a=\frac{\lambda_2+\sqrt{\lambda_2^2-4(\alpha+3\lambda_1\lambda_2-3\lambda_1^2)}}{2}$, using the equation (52) again, we get $a-\frac{\alpha-\lambda_2^2+3\lambda_1\lambda_2}{\lambda_2}=\frac{p_1\alpha_2}{-\lambda_2p_1}=-\frac{\alpha_2}{\lambda_2}$, so $\alpha_2=\alpha-\lambda_2^2+3\lambda_1\lambda_2-\lambda_2a$. By the equation $(44b)$, we get $(3\lambda_1-\lambda_2)^2\alpha=18\lambda_1^4-6\lambda_1\lambda_2^3+24\lambda_1^2\lambda_2^2-36\lambda_1^3\lambda_2$, if $3\lambda_1=\lambda_2$, then $0=-36\lambda_1^2$, which is a contradiction, so $3\lambda_1\neq\lambda_2$ and $\alpha=\frac{18\lambda_1^4-6\lambda_1\lambda_2^3+24\lambda_1^2\lambda_2^2-36\lambda_1^3\lambda_2}{(3\lambda_1-\lambda_2)^2}$, so $a=\frac{\lambda_2+\sqrt{\frac{(6\lambda_1^2-3\lambda_1\lambda_2-\lambda_2^2)^2}{(3\lambda_1-\lambda_2)^2}}}{2}$.

If $\frac{6\lambda_1^2-3\lambda_1\lambda_2-\lambda_2^2}{3\lambda_1-\lambda_2}>0$, then $a=\frac{3\lambda_1^2-\lambda_2^2}{3\lambda_1-\lambda_2},\ \alpha_2=\frac{18\lambda_1^4-2\lambda_2^4+6\lambda_1\lambda_2^3-18\lambda_1^3\lambda_2}{(3\lambda_1-\lambda_2)^2}=\frac{(3\lambda_1^2-\lambda_2^2)(3\lambda_1^2-3\lambda_1\lambda_2+\lambda_2^2)}{(3\lambda_1-\lambda_2)^2}$, since $3\lambda_1^2-3\lambda_1\lambda_2+\lambda_2^2=3\big(\lambda_1-\frac{1}{2}\lambda_2\big)^2+\frac{1}{4}\lambda_2^2\neq 0$, and $\alpha_2\neq 0$, we have $3\lambda_1^2\neq\lambda_2^2$, so $a\neq 0$.

If $\frac{6\lambda_1^2-3\lambda_1\lambda_2-\lambda_2^2}{3\lambda_1-\lambda_2}<0$, then $a=\frac{3\lambda_1\lambda_2-\lambda_2^2}{3\lambda_1-\lambda_2},\ \alpha_2=\frac{18\lambda_1^4-\lambda_2^4+6\lambda_1\lambda_2^3-15\lambda_1^2\lambda_2^2}{(3\lambda_1-\lambda_2)^2}\neq 0$, so $18\lambda_1^4-\lambda_2^4+6\lambda_1\lambda_2^3-15\lambda_1^2\lambda_2^2\neq 0$.

Hence, we have $\underline{\lambda_2\neq 3\lambda_1,\ \lambda_2^2\neq 3\lambda_1^2,\ \frac{6\lambda_1^2-3\lambda_1\lambda_2-\lambda_2^2}{3\lambda_1-\lambda_2}>0,\ p_1\neq 0,\ p_2=0,\ \alpha_1=0,\ \alpha=\frac{18\lambda_1^4-6\lambda_1\lambda_2^3+24\lambda_1^2\lambda_2^2-36\lambda_1^3\lambda_2}{(3\lambda_1-\lambda_2)^2},\ \alpha_2=\frac{18\lambda_1^4-2\lambda_2^4+6\lambda_1\lambda_2^3-18\lambda_1^3\lambda_2}{(3\lambda_1-\lambda_2)^2},\ \phi=ce^{\frac{3\lambda_1^2-\lambda_2^2}{(3\lambda_1-\lambda_2)p_1}t}.}$
$\underline{\text{or } \lambda_2\neq 3\lambda_1,\ 18\lambda_1^4-\lambda_2^4+6\lambda_1\lambda_2^3-15\lambda_1^2\lambda_2^2\neq 0,\ \frac{6\lambda_1^2-3\lambda_1\lambda_2-\lambda_2^2}{3\lambda_1-\lambda_2}<0,\ p_1\neq 0,\ p_2=0,\ \alpha_1=0,\alpha=\frac{18\lambda_1^4-6\lambda_1\lambda_2^3+24\lambda_1^2\lambda_2^2-36\lambda_1^3\lambda_2}{(3\lambda_1-\lambda_2)^2},\ \alpha_2=\frac{18\lambda_1^4-\lambda_2^4+6\lambda_1\lambda_2^3-15\lambda_1^2\lambda_2^2}{(3\lambda_1-\lambda_2)^2},\ \phi=ce^{\frac{3\lambda_1\lambda_2-3\lambda_1^2}{(3\lambda_1-\lambda_2)p_1}t}.}$
Especially when $\lambda_1=\lambda_2<0$, we have $p_1\neq 0,\ p_2=0,\ \alpha=\alpha_1=0,\ \alpha_2=2\lambda_1^2,\ \phi=c$.
$2'$. $\underline{p_1\alpha_2=0.}$ 1°. $\underline{p_1=0.}$ Then $\zeta=2p_2,\ \eta=2p_2^2,\ \frac{\eta}{\zeta^2}=\frac{1}{2}$. By the equation (52), we get $a=\frac{\alpha-\lambda_2^2+3\lambda_1\lambda_2}{2\lambda_2}$, on the other hand, $a=\frac{\lambda_2+\sqrt{\lambda_2^2-2(\alpha+3\lambda_1\lambda_2-3\lambda_1^2)}}{2}$, so we get

$$\alpha^2+(6\lambda_1\lambda_2-2\lambda_2^2)\alpha+3\lambda_1^2\lambda_2^2-6\lambda_1^3\lambda_2+3\lambda_1^4=0.\tag{53}$$

when $\lambda_1=\lambda_2$, we get $\alpha^2+4\lambda_1^2\alpha=0$, then $\alpha=0$ or $\alpha=-4\lambda_1^2$. If $\alpha=0$, we have

$a=\lambda_1=\lambda_2>0$, by the equation $(44c)$, we get $\alpha_2=0,\ \phi=\psi^{\frac{\zeta}{\eta}}=c_0e^{\frac{\lambda_1}{p_2}t}$. If $\alpha=-4\lambda_1^2$, then $\lambda_1=\lambda_2<0,\ a=-\lambda_1$, then by the equation $(44c)$, we get $\lambda_1=0$, which is a contradiction.

When $\lambda_1\neq\lambda_2,\ \Delta=8\lambda_2^2(3\lambda_1^2-\lambda_2^2)$, if $\lambda_2^2>3\lambda_1^2$, we have no solution; if $\lambda_2^2=3\lambda_1^2$, we have $\underline{a=0,\ \alpha_1=\alpha_2=0,\ \alpha=\lambda_2^2-3\lambda_1\lambda_2,\ \phi=c;}$ if $\lambda_2^2<3\lambda_1^2$, we have $\alpha=\lambda_2^2-3\lambda_1\lambda_2+\sqrt{2\lambda_2^2(3\lambda_1^2-\lambda_2^2)}$ or $\alpha=\lambda_2^2-3\lambda_1\lambda_2-\sqrt{2\lambda_2^2(3\lambda_1^2-\lambda_2^2)}$. By the equation $(44c)$, we have $\alpha_2=0$ and

$$-2a^2+(3\lambda_1+\lambda_2)a=\alpha-\lambda_2^2+3\lambda_1\lambda_2. \tag{54}$$

then we have $\lambda_1=\lambda_2>0$ or $\lambda_1=\lambda_2<0$, but when $\lambda_1=\lambda_2<0$, it is not satisfies the equation $(44c)$, so we get $\underline{\lambda_1=\lambda_2>0,\ p_1=0,\ p_2\neq0,\ \alpha=\alpha_1=\alpha_2=0,\ \phi=ce^{\frac{\lambda_1}{p_2}t}}$.

2°. $\underline{\alpha_2=0.}$ Since $p_1=0$ we have discussed, so we assume that $p_1\neq0$. By the equation (52), we have $a=\frac{\alpha-\lambda_2^2+3\lambda_1\lambda_2}{\lambda_2}\frac{\eta}{\zeta^2},\ \phi^\zeta=ce^{a\frac{\zeta^2}{\eta}t}$, then by the equation$(44b)$, we have $\alpha(\alpha-\lambda_2^2+3\lambda_1\lambda_2)=0$. So when $\lambda_2=3\lambda_1$, we have $\alpha=0$; when $\lambda_2\neq3\lambda_1$, we have $\alpha=0$ or $\alpha=\lambda_2^2-3\lambda_1\lambda_2\neq0$.

When $\lambda_2=3\lambda_1,\ \alpha=0$, then $a=0,\ \eta=0$, which is a contradiction.

When $\lambda_2\neq3\lambda_1,\ \alpha=\lambda_2^2-3\lambda_1\lambda_2\neq0$, then $a=0$ and $(\lambda_2^2-3\lambda_1^2)\eta=0$.

If $\lambda_2^2\neq3\lambda_1^2$, then $\eta=0$, which is a contradiction;

If $\lambda_2^2=3\lambda_1^2$, which satisfies $\lambda_2\neq3\lambda_1$, then $\phi^\zeta=c$, which satisfies equations $(44a)-(44c)$. so we have $\underline{\lambda_2^2=3\lambda_1^2,\ p_1\neq0,\ p_2\neq-\frac{1}{2}p_1.\ \alpha=\lambda_2^2-3\lambda_1\lambda_2,\ \alpha_1=\alpha_2=0,\ \phi=c.}$

When $\lambda_2\neq3\lambda_1,\ \alpha=0$, then $a=(3\lambda_1-\lambda_2)\frac{\eta}{\zeta^2}$, by the equation $(44a)$, we get $\frac{\eta}{\zeta^2}=\frac{3\lambda_1^2-\lambda_2^2}{(3\lambda_1-\lambda_2)^2}>0$ and $\lambda_2^2<3\lambda_1^2$, then $a=\frac{3\lambda_1^2-\lambda_2^2}{3\lambda_1-\lambda_2}$. Since $a>\frac{\lambda_2}{2}$, we get $\frac{6\lambda_1^2-3\lambda_1\lambda_2-\lambda_2^2}{3\lambda_1-\lambda_2}>0$, considering that$\frac{p_1^2+2p_2^2}{(p_1+2p_2)^2}=\frac{\eta}{\zeta^2}=\frac{3\lambda_1^2-\lambda_2^2}{(3\lambda_1-\lambda_2)^2}$, we have $(3\lambda_1^2-3\lambda_1\lambda_2+\lambda_2^2)p_1^2-(6\lambda_1^2-2\lambda_2^2)p_1p_2=(3\lambda_1^2-6\lambda_1\lambda_2+3\lambda_2^2)p_2^2$, no matter $\lambda_1=\lambda_2$ or $\lambda_1\neq\lambda_2$, we can get $p_2\neq0$.

So we get $\underline{\lambda_2\neq3\lambda_1,\ \lambda_2^2<3\lambda_1^2,\ \frac{6\lambda_1^2-3\lambda_1\lambda_2-\lambda_2^2}{3\lambda_1-\lambda_2}>0,\ \frac{\eta}{\zeta^2}=\frac{3\lambda_1^2-\lambda_2^2}{(3\lambda_1-\lambda_2)^2},\ \alpha=\alpha_1=\alpha_2=0,}$
$\underline{\phi=ce^{\frac{3\lambda_1^2-\lambda_2^2}{3\lambda_1-\lambda_2}\frac{\zeta}{\eta}t}.}$

3) $\underline{c_1\neq0,\ c_2\neq0,\ b\neq0.}$

$1'$. $\underline{p_2\neq0.}$ Then e^{at},e^{bt} and $(c_1e^{at}+c_2e^{bt})^{1-2p_2\frac{\zeta}{\eta}}$ are linearly independent, by the equation (46), we have $\left[a-\frac{\alpha-\lambda_2^2+3\lambda_1\lambda_2}{\lambda_2}\frac{\eta}{\zeta^2}\right]c_1=0,\quad\left[b-\frac{\alpha-\lambda_2^2+3\lambda_1\lambda_2}{\lambda_2}\frac{\eta}{\zeta^2}\right]c_2=0.$

Considering that $c_1\neq0,\ c_2\neq0$, we have $a=b=\frac{\alpha-\lambda_2^2+3\lambda_1\lambda_2}{\lambda_2}\frac{\eta}{\zeta^2}$, which is a contradiction.

$2'$. $\underline{p_2=0.}$ Then by the equation (46), we have

$a-\frac{\alpha-\lambda_2^2+3\lambda_1\lambda_2}{\lambda_2}\frac{\eta}{\zeta^2}-\frac{p_1\alpha_2}{\lambda_2(p_2-p_1)}\frac{\eta}{\zeta^2}=0, \qquad b-\frac{\alpha-\lambda_2^2+3\lambda_1\lambda_2}{\lambda_2}\frac{\eta}{\zeta^2}-\frac{p_1\alpha_2}{\lambda_2(p_2-p_1)}\frac{\eta}{\zeta^2}=0.$
Then $a=b$, which is a contradiction.

4) $\underline{c_1\neq 0,\ c_2\neq 0,\ b=0.}$ Then $a=\lambda_2\neq 0$ and

$$ac_1e^{at}=\frac{p_1\alpha_2}{\lambda_2(p_2-p_1)}\frac{\eta}{\zeta^2}(c_1e^{at}+c_2)^{1-2p_2\frac{\zeta}{\eta}}+\frac{\alpha-\lambda_2^2+3\lambda_1\lambda_2}{\lambda_2}\frac{\eta}{\zeta^2}(c_1e^{at}+c_2). \tag{55}$$

$1'.\ \underline{1-2p_2\frac{\zeta}{\eta}\neq 0.}$

If $p_2\neq 0$, then e^{at} and $(c_1e^{at}+c_2)^{1-2p_2\frac{\zeta}{\eta}}$ are linearly independent, by the equation (55), we have $a=0$, this is a contradiction with $a=\lambda_2\neq 0$.

If $p_2=0$, using the same method we can get $a=0$, this is a contradiction with $a=\lambda_2\neq 0$.

$2'.\ \underline{1-2p_2\frac{\zeta}{\eta}=0.}$

Then $p_2\neq 0,\ \eta=2p_2\zeta$ and the equation (55) becomes

$$\Big(\lambda_2-\frac{\alpha-\lambda_2^2+3\lambda_1\lambda_2}{\lambda_2}\frac{\eta}{\zeta^2}\Big)c_1e^{at}-\frac{p_1\alpha_2}{\lambda_2(p_2-p_1)}\frac{\eta}{\zeta^2}-\frac{\alpha-\lambda_2^2+3\lambda_1\lambda_2}{\lambda_2}\frac{\eta}{\zeta^2}=0. \tag{56}$$

Then

$$\lambda_2=\frac{\alpha-\lambda_2^2+3\lambda_1\lambda_2}{\lambda_2}\frac{\eta}{\zeta^2}. \tag{57}$$

Since $b=0$ which means $\lambda_2=\sqrt{\lambda_2^2-4(\alpha+3\lambda_1\lambda_2-3\lambda_1^2)\frac{\eta}{\zeta^2}}$, we get $\alpha=3\lambda_1^2-3\lambda_1\lambda_2$, then using the equation (57), we have $\lambda_2=\frac{3\lambda_1^2-\lambda_2^2}{\lambda_2}\frac{\eta}{\zeta^2}$ and $\lambda_2^2\neq 3\lambda_1^2$, considering that $\eta=2p_2\zeta$, we get $p_1=\frac{6\lambda_1^2-4\lambda_2^2}{\lambda_2^2}p_2$.

If $3\lambda_1^2=2\lambda_2^2$, then $p_1=0$, by $\eta=2p_2\zeta$ we have $p_2=0$, which is a contradiction.

If $3\lambda_1^2\neq 2\lambda_2^2$, then $p_1\neq 0$, and by $\eta=2p_2\zeta$ we have $18\lambda_1^4-30\lambda_1^2\lambda_2^2+11\lambda_2^4=0$, then $\lambda_1^2=\frac{5\pm\sqrt{3}}{6}\lambda_2^2$, which satisfies $\lambda_2^2\neq 3\lambda_1^2$ and $3\lambda_1^2\neq 2\lambda_2^2$. Then $\alpha_2=\frac{(p_2-p_1)(\lambda_2^2-3\lambda_1\lambda_2)}{p_1}$, $p_1=(1\pm\sqrt{3})p_2\neq 0,\ \zeta=(3\pm\sqrt{3})p_2,\ \eta=(6\pm 2\sqrt{3})p_2^2,\ \phi=ce^{\frac{\lambda_2}{2p_1}t}$.

So we get $\underline{\lambda_1^2=\frac{5\pm\sqrt{3}}{6}\lambda_2^2,\ p_1=(1\pm\sqrt{3})p_2\neq 0,\ \alpha=3\lambda_1^2-3\lambda_1\lambda_2,\ \alpha_1=0,\ \alpha_2=\frac{(p_2-p_1)(\lambda_2^2-3\lambda_1\lambda_2)}{p_1},\ \phi=ce^{\frac{\lambda_2}{2p_1}t}.}$

$\alpha=\frac{\lambda_2^2\zeta^2}{4\eta}+3\lambda_1^2-3\lambda_1\lambda_2,\ \psi=c_1e^{\frac{\lambda_2}{2}t}+c_2te^{\frac{\lambda_2}{2}t}$. By the equation (45), we have

$$\Big[\frac{\lambda_2}{2}c_1+c_2-a_0c_1+\Big(\frac{\lambda_2}{2}c_2-a_0c_2\Big)t\Big]e^{\frac{\lambda_2}{2}t}=\frac{p_1\alpha_2}{\lambda_2(p_2-p_1)}\frac{\eta}{\zeta^2}(c_1+c_2t)^{1-2p_2\frac{\zeta}{\eta}}(e^{\frac{\lambda_2}{2}t})^{1-2p_2\frac{\zeta}{\eta}}, \tag{58}$$

where $a_0=\frac{\lambda_2}{4}+\frac{3\lambda_1^2-\lambda_2^2}{\lambda_2^2}\frac{\eta}{\zeta^2}$.

1) $\underline{c_2\neq 0.}$ Then by the equation (58), we have $p_2=0$, by the equation (44c), we

get $\alpha_2+\lambda_2\frac{(\phi^\zeta)'}{\phi^\zeta}=\alpha-\lambda_2^2+3\lambda_1\lambda_2$, then $\phi^\zeta=c_0e^{\frac{\alpha-\lambda_2^2+3\lambda_1\lambda_2-\alpha_2}{\lambda_2}t}$; on the other hand, $\phi^\zeta=\psi^{\frac{\zeta^2}{\eta}}=\Big(c_1e^{\frac{\lambda_2}{2}t}+c_2te^{\frac{\lambda_2}{2}t}\Big)^{\frac{\zeta^2}{\eta}}$, then $c_0e^{\frac{\alpha-\lambda_2^2+3\lambda_1\lambda_2-\alpha_2}{\lambda_2}t}=\Big(c_1e^{\frac{\lambda_2}{2}t}+c_2te^{\frac{\lambda_2}{2}t}\Big)^{\frac{\zeta^2}{\eta}}$, this is a contradiction with $c_2\neq 0$.

2) $\underline{c_2=0.}$ Then the equation (58) becomes

$$\Big(\frac{\lambda_2}{2}c_1-a_0c_1\Big)e^{\frac{\lambda_2}{2}t}=\frac{p_1\alpha_2}{\lambda_2(p_2-p_1)}\frac{\eta}{\zeta^2}c_1^{1-2p_2\frac{\zeta}{\eta}}(e^{\frac{\lambda_2}{2}t})^{1-2p_2\frac{\zeta}{\eta}}. \tag{59}$$

1′. $\underline{a_0=\frac{\lambda_2}{2}}$. By the equation (59), we get $p_1\alpha_2=0$. Considering that $a_0=\frac{\lambda_2}{4}+\frac{3\lambda_1^2-\lambda_2^2}{\lambda_2^2}\frac{\eta}{\zeta^2}$, we have $\lambda_2^2\neq 3\lambda_1^2$ and $\frac{\lambda_2^2}{4}\frac{\zeta^2}{\eta}=3\lambda_1^2-\lambda_2^2$, then $\alpha=6\lambda_1^2-3\lambda_1\lambda_2-\lambda_2^2$.

If $p_1=0$, then $\zeta=2p_2,\ \eta=2p_2^2,\ \frac{\eta}{\zeta^2}=\frac{1}{2}$. By $\frac{\lambda_2^2}{4}\frac{\zeta^2}{\eta}=3\lambda_1^2-\lambda_2^2$, we have $\lambda_2^2=2\lambda_1^2$ and $\alpha=2\lambda_2^2-3\lambda_1\lambda_2$. By the equation (44$b$), we get $\phi^\zeta=c_0e^{\lambda_2t}$, then by the equation (44c), we get $\lambda_2=\frac{3}{2}\lambda_1$, this is a contradiction with $\lambda_2^2=2\lambda_1^2$.

If $p_1\neq 0,\ \alpha_2=0$, then $\psi=c_1e^{\frac{\lambda_2}{2}t},\ \phi^\zeta=c_0e^{\frac{2(3\lambda_1^2-\lambda_2^2)}{\lambda_2}t}$, then by the equation (44b), we have $\alpha=6\lambda_1^2-3\lambda_1\lambda_2-\lambda_2^2=0$, which satisfies the equation (44c). $\phi=c_0e^{\frac{\lambda_2}{2}\frac{\zeta}{\eta}t}$.

So we get $\underline{p_1\neq 0,\ \alpha=6\lambda_1^2-3\lambda_1\lambda_2-\lambda_2^2=0,\ \alpha_1=\alpha_2=0,\ \phi=c_0e^{\frac{\lambda_2}{2}\frac{\zeta}{\eta}t}.}$

2′. $\underline{a_0\neq\frac{\lambda_2}{2}}$. By the equation (59), we have $p_2=0$, then $\zeta=p_1,\ \eta=p_1^2,\ \frac{\zeta^2}{\eta}=1$, so $\alpha=\frac{\lambda_2^2}{4}+3\lambda_1^2-3\lambda_1\lambda_2,\ \phi^\zeta=c_1e^{\frac{\lambda_2}{2}t}$. By equation (44$b$), we have $6\lambda_1^2-3\lambda_1\lambda_2-\lambda_2^2=0$, by the equation (44$c$), we have $\alpha_2=3\lambda_1^2-\frac{5}{4}\lambda_2^2$.

So we get $\underline{6\lambda_1^2-3\lambda_1\lambda_2-\lambda_2^2=0,\ p_1\neq 0,\ p_2=0,\ \alpha=\frac{\lambda_2^2}{4}+3\lambda_1^2-3\lambda_1\lambda_2,\ \alpha_2=3\lambda_1^2-\frac{5}{4}\lambda_2^2,\ \phi=ce^{\frac{\lambda_2}{2\zeta}t}.}$

$\alpha>\frac{\lambda_2^2\zeta^2}{4\eta}+3\lambda_1^2-3\lambda_1\lambda_2,\ \psi=c_1e^{\frac{\lambda_2}{2}t}cos(at)+c_2e^{\frac{\lambda_2}{2}t}sin(at),\ a=\frac{\sqrt{4(\alpha+3\lambda_1\lambda_2-3\lambda_1^2)\frac{\eta}{\zeta^2}-\lambda_2^2}}{2}$.

By the equation (45), we have

$$\begin{aligned}&\Big(\frac{\lambda_2}{2}c_1+ac_2\Big)cos(at)+\Big(-ac_1+\frac{\lambda_2}{2}c_2\Big)sin(at)\\ =\ &\frac{p_1\alpha_2}{\lambda_2(p_2-p_1)}\frac{\eta}{\zeta^2}\big(c_1cos(at)+c_2sin(at)\big)^{1-2p_2\frac{\zeta}{\eta}}e^{-\lambda_2p_2\frac{\zeta}{\eta}t}\\ +\ &\frac{\alpha-\lambda_2^2+3\lambda_1\lambda_2}{\lambda_2}\frac{\eta}{\zeta^2}\big(c_1cos(at)+c_2sin(at)\big).\end{aligned} \tag{60}$$

1) $\underline{p_2\neq 0.}$ Then by the equation (60), we get $p_1\alpha_2=0$ and

$$\frac{\lambda_2}{2}c_1+ac_2=\frac{\alpha-\lambda_2^2+3\lambda_1\lambda_2}{\lambda_2}\frac{\eta}{\zeta^2}c_1; \tag{61a}$$

$$-ac_1+\frac{\lambda_2}{2}c_2=\frac{\alpha-\lambda_2^2+3\lambda_1\lambda_2}{\lambda_2}\frac{\eta}{\zeta^2}c_2. \tag{61b}$$

$(61a)\times c_2-(61b)\times c_1$, we get $c_1^2+c_2^2=0$, this is a contradiction.
2) $\underline{p_2=0.}$ Then by the equation (60), we have

$$\frac{\lambda_2}{2}c_1+ac_2=\frac{p_1\alpha_2}{\lambda_2(p_2-p_1)}\frac{\eta}{\zeta^2}c_1+\frac{\alpha-\lambda_2^2+3\lambda_1\lambda_2}{\lambda_2}\frac{\eta}{\zeta^2}c_1; \tag{62a}$$

$$-ac_1+\frac{\lambda_2}{2}c_2=\frac{p_1\alpha_2}{\lambda_2(p_2-p_1)}\frac{\eta}{\zeta^2}c_2+\frac{\alpha-\lambda_2^2+3\lambda_1\lambda_2}{\lambda_2}\frac{\eta}{\zeta^2}c_2. \tag{62b}$$

$(62a)\times c_2-(62b)\times c_1$, we get $c_1^2+c_2^2=0$, this is a contradiction.
So we have no solution in .

According to above discussions, we get the following Theorem:

Theorem 4.19. *Let $M=I\times_{\phi^{p_1}}F_1\times_{\phi^{p_2}}F_2\cdots\times_{\phi^{p_m}}F_m$ be a generalized Kasner space-time,* $\dim F_1=1,\ \dim F_2=2$ *and $P=\frac{\partial}{\partial t}$. Then $(M,\overline{\nabla})$ is Einstein with the Einstein constant α if and only if one of the following conditions is satisfied:*
$(1)\lambda_2^2=3\lambda_1^2,\ p_1=p_2=0,\ \alpha=3\lambda_1^2-3\lambda_1\lambda_2=\lambda_2^2-3\lambda_1\lambda_2,\ \alpha_1=\alpha_2=0;$
$(2)\lambda_2^2=3\lambda_1^2,\ p_1^2+p_2^2\neq 0,\ \alpha=3\lambda_1^2-3\lambda_1\lambda_2=\lambda_2^2-3\lambda_1\lambda_2,\ \alpha_1=\alpha_2=0,\ \phi=c;$
$(3)\lambda_2^2\neq 3\lambda_1^2,\ \lambda_2\neq 3\lambda_1,\ \frac{6\lambda_1^2-3\lambda_1\lambda_2-\lambda_2^2}{3\lambda_1-\lambda_2}\neq 0,\ p_1\neq 0,\ p_2=0,\ \alpha=\frac{18\lambda_1^4-6\lambda_1\lambda_2^3+24\lambda_1^2\lambda_2^2-36\lambda_1^3\lambda_2}{(3\lambda_1-\lambda_2)^2},$
$\alpha_1=0,\alpha_2=\frac{18\lambda_1^4-2\lambda_2^4+6\lambda_1\lambda_2^3-18\lambda_1^3\lambda_2}{(3\lambda_1-\lambda_2)^2},\ \phi=ce^{\frac{3\lambda_1^2-\lambda_2^2}{(3\lambda_1-\lambda_2)p_1}t};$
$(4)\lambda_2^2<3\lambda_1^2,\ \lambda_2\neq 3\lambda_1,\ \frac{6\lambda_1^2-3\lambda_1\lambda_2-\lambda_2^2}{3\lambda_1-\lambda_2}\neq 0,\ p_1\neq 0,\ p_2\neq-\frac{1}{2}p_1,\ \frac{\eta}{\zeta^2}=\frac{3\lambda_1^2-\lambda_2^2}{(3\lambda_1-\lambda_2)^2},\ \alpha=$
$\alpha_1=\alpha_2=0,\phi=ce^{\frac{3\lambda_1-\lambda_2}{\zeta}t};$
$(5)\lambda_2\neq 3\lambda_1,\frac{6\lambda_1^2-3\lambda_1\lambda_2-\lambda_2^2}{3\lambda_1-\lambda_2}<0,18\lambda_1^4-\lambda_2^4+6\lambda_1\lambda_2^3-15\lambda_1^2\lambda_2^2\neq 0,p_1\neq 0,p_2=0,\alpha_1=$
$0,\alpha=\frac{18\lambda_1^4-6\lambda_1\lambda_2^3+24\lambda_1^2\lambda_2^2-36\lambda_1^3\lambda_2}{(3\lambda_1-\lambda_2)^2},\ \alpha_2=\frac{18\lambda_1^4-\lambda_2^4+6\lambda_1\lambda_2^3-15\lambda_1^2\lambda_2^2}{(3\lambda_1-\lambda_2)^2},\ \phi=ce^{\frac{3\lambda_1\lambda_2-3\lambda_1^2}{(3\lambda_1-\lambda_2)p_1}t};$
$(6)\lambda_1=\lambda_2,\ p_1=0,\ p_2\neq 0,\ \alpha=\alpha_1=\alpha_2=0,\ \phi=c_0e^{\frac{\lambda_1}{p_2}t};$
$(7)\lambda_1^2=\frac{5\pm\sqrt{3}}{6}\lambda_2^2,\ p_1=(1\pm\sqrt{3})p_2\neq 0,\ \alpha=3\lambda_1^2-3\lambda_1\lambda_2,\ \alpha_1=0,\ \alpha_2=\frac{(p_2-p_1)(\lambda_2^2-3\lambda_1\lambda_2)}{p_1},$
$\phi=ce^{\frac{\lambda_2}{2p_1}t};$
$(8)6\lambda_1^2-3\lambda_1\lambda_2-\lambda_2^2=0,\ p_1\neq 0,\ \alpha=6\lambda_1^2-3\lambda_1\lambda_2-\lambda_2^2=0,\ \alpha_1=\alpha_2=0,\ \phi=ce^{\frac{\lambda_2}{2}\frac{\zeta}{\eta}t};$
$(9)6\lambda_1^2-3\lambda_1\lambda_2-\lambda_2^2=0,\ p_1\neq 0,\ p_2=0,\ \frac{\zeta^2}{\eta}=1,\ \alpha=\frac{\lambda_2^2}{4}+3\lambda_1^2-3\lambda_1\lambda_2,\ \alpha_2=$
$3\lambda_1^2-\frac{5}{4}\lambda_2^2,\phi=ce^{\frac{\lambda_2}{2}\frac{\zeta}{\eta}t}.$

4.3.4 Type (II) generalized Kasner space-times with a quarter-symmetric connection with constant scalar curvature

By Proposition 4.17, then (F_2,∇^{F_2}) has constant scalar curvature S^{F_2} and

$$\overline{S}=\frac{S^{F_2}}{\phi^{2p_2}}-2\zeta\frac{\phi''}{\phi}-(\eta+\zeta^2-2\zeta)\frac{(\phi')^2}{\phi^2}+3(\lambda_1+\lambda_2)\zeta\frac{\phi'}{\phi}+3(\lambda_1^2+\lambda_2^2-4\lambda_1\lambda_2). \tag{63}$$

$\zeta = 0$. (1) $\eta = 0$. then $p_1 = p_2 = 0$ and $\overline{S} = S^{F_2} + 3(\lambda_1^2 + \lambda_2^2 - 4\lambda_1\lambda_2)$;
(2) $\eta \neq 0$. then $\overline{S} = \frac{S^{F_2}}{\phi^{2p_2}} - \eta\frac{(\phi')^2}{\phi^2} + 3(\lambda_1^2 + \lambda_2^2 - 4\lambda_1\lambda_2)$, which means

$$\eta\frac{(\phi')^2}{\phi^2} = \frac{S^{F_2}}{\phi^{2p_2}} - [\overline{S} - 3(\lambda_1^2 + \lambda_2^2 - 4\lambda_1\lambda_2)]. \tag{64}$$

$\zeta \neq 0$. Putting $\phi = \psi^{\frac{2\zeta}{\eta+\zeta^2}}$, we get

$$-\frac{4\zeta^2}{\eta+\zeta^2}\psi'' + \frac{6(\lambda_1+\lambda_2)\zeta^2}{\eta+\zeta^2}\psi' + (3\lambda_1^2 + 3\lambda_2^2 - 12\lambda_1\lambda_2 - \overline{S})\psi + S^{F_2}\psi^{1-\frac{4p_2\zeta}{\eta+\zeta^2}} = 0. \tag{65}$$

4.3.5 Classification of Einstein Type (III) generalized Kasner space-times with a quarter-symmetric connection

Considering $\dim F_1 = \dim F_2 = \dim F_3 = 1$, by Remark 4, we get $\alpha_i = 0$ and by Proposition 4.16, we have

$$\zeta\Big(\lambda_2\frac{\phi'}{\phi} - \frac{\phi''}{\phi}\Big) - (\eta-\zeta)\frac{(\phi')^2}{\phi^2} + 3(\lambda_1^2 - \lambda_1\lambda_2) = \alpha; \tag{66a}$$

$$-p_1\Big[\frac{\phi''}{\phi} + (\zeta-1)\frac{(\phi')^2}{\phi^2} + (\lambda_2 - 3\lambda_1)\frac{\phi'}{\phi}\Big] + \lambda_2\zeta\frac{\phi'}{\phi} = \alpha - \lambda_2^2 + 3\lambda_1\lambda_2. \tag{66b}$$

$$-p_2\Big[\frac{\phi''}{\phi} + (\zeta-1)\frac{(\phi')^2}{\phi^2} + (\lambda_2 - 3\lambda_1)\frac{\phi'}{\phi}\Big] + \lambda_2\zeta\frac{\phi'}{\phi} = \alpha - \lambda_2^2 + 3\lambda_1\lambda_2. \tag{66c}$$

$$-p_3\Big[\frac{\phi''}{\phi} + (\zeta-1)\frac{(\phi')^2}{\phi^2} + (\lambda_2 - 3\lambda_1)\frac{\phi'}{\phi}\Big] + \lambda_2\zeta\frac{\phi'}{\phi} = \alpha - \lambda_2^2 + 3\lambda_1\lambda_2. \tag{66d}$$

$\zeta = \eta = 0$. By the equation (66a), we have $\alpha = 3\lambda_1^2 - 3\lambda_1\lambda_2$, and by the equation (66b), we have $\alpha = \lambda_2^2 - 3\lambda_1\lambda_2$, then we get $\lambda_2^2 = 3\lambda_1^2$.
So we obtain $\underline{\lambda_2^2 = 3\lambda_1^2,\ \alpha = 3\lambda_1^2 - 3\lambda_1\lambda_2 = \lambda_2^2 - 3\lambda_1\lambda_2,\ \alpha_i = 0,\ \zeta = \eta = 0.}$
$\zeta = 0,\ \eta \neq 0$. (66b) + (66c) + (66d), we get $\alpha = \lambda_2^2 - 3\lambda_1\lambda_2$; by the equation (66$a$), we have $\frac{(\phi')^2}{\phi^2} = \frac{3\lambda_1^2 - \lambda_2^2}{\eta}$.
1) $3\lambda_1^2 - \lambda_2^2 < 0$, we have no solution.
2) $3\lambda_1^2 - \lambda_2^2 = 0$, then $\phi = c$, which satisfies the equation (66a).
3) $3\lambda_1^2 - \lambda_2^2 > 0$, then $\phi = c_0 e^{\pm\sqrt{\frac{3\lambda_1^2-\lambda_2^2}{\eta}}t}$, since $\eta \neq 0$, so at least one $p_i \neq 0$, we assume $p_1 \neq 0$, by the equation (66b), we get $\lambda_2 = 3\lambda_1$, but by $3\lambda_1^2 - \lambda_2^2 > 0$, we get $\lambda_1^2 < 0$, which is a contradiction.
So we have $\underline{\lambda_2^2 = 3\lambda_1^2,\ \alpha = 3\lambda_1^2 - 3\lambda_1\lambda_2 = \lambda_2^2 - 3\lambda_1\lambda_2,\ \alpha_i = 0,\ \zeta = 0,\ \eta \neq 0,\ \phi = c}$ in case .

$\zeta \neq 0$, then $\eta \neq 0$. If $p_1 = p_2 = p_3$, we get type (I), if $p_1 = p_2$ or $p_2 = p_3$ or $p_1 = p_3$, we get type (II), so $p_1 \neq p_2 \neq p_3$. Let $\phi = \psi^{\frac{\zeta}{\eta}}$, then equations $(66a) - (66d)$ becomes

$$\frac{\zeta^2}{\eta}\frac{\lambda_2\psi' - \psi''}{\psi} = \alpha + 3\lambda_1\lambda_2 - 3\lambda_2^2 \tag{67a}$$

$$\frac{p_1}{\zeta}\Big[-\frac{(\phi^\zeta)''}{\phi^\zeta} + (3\lambda_1 - \lambda_2)\frac{(\phi^\zeta)'}{\phi^\zeta}\Big] + \lambda_2\frac{(\phi^\zeta)'}{\phi^\zeta} = \alpha - \lambda_2^2 + 3\lambda_1\lambda_2 \tag{67b}$$

$$\frac{p_2}{\zeta}\Big[-\frac{(\phi^\zeta)''}{\phi^\zeta} + (3\lambda_1 - \lambda_2)\frac{(\phi^\zeta)'}{\phi^\zeta}\Big] + \lambda_2\frac{(\phi^\zeta)'}{\phi^\zeta} = \alpha - \lambda_2^2 + 3\lambda_1\lambda_2 \tag{67c}$$

$$\frac{p_3}{\zeta}\Big[-\frac{(\phi^\zeta)''}{\phi^\zeta} + (3\lambda_1 - \lambda_2)\frac{(\phi^\zeta)'}{\phi^\zeta}\Big] + \lambda_2\frac{(\phi^\zeta)'}{\phi^\zeta} = \alpha - \lambda_2^2 + 3\lambda_1\lambda_2 \tag{67d}$$

$(67b) \times p_2 - (67c) \times p_1$ and considering that $p_1 \neq p_2$, we get

$$\lambda_2\frac{(\phi^\zeta)'}{\phi^\zeta} = \alpha - \lambda_2^2 + 3\lambda_1\lambda_2 \tag{68a}$$

$$\frac{(\phi^\zeta)'}{\phi^\zeta} = \frac{\alpha}{\lambda_2} - \lambda_2 + 3\lambda_1 \tag{68b}$$

by equations $(67b)$ and $(68a)$, we get $-\frac{(\phi^\zeta)''}{\phi^\zeta} + (3\lambda_1 - \lambda_2)\frac{(\phi^\zeta)'}{\phi^\zeta} = 0$, then by the equation $(68b)$, we have

$$\frac{(\phi^\zeta)''}{\phi^\zeta} = (\frac{3\lambda_1}{\lambda_2} - 1)\alpha + 9\lambda_1^2 - 6\lambda_1\lambda_2 + \lambda_2^2 \tag{69}$$

on the other hand, using the equation $(68b)$, we get

$$\phi^\zeta = c_0 e^{(\frac{\alpha}{\lambda_2} - \lambda_2 + 3\lambda_1)t} \tag{70}$$

by equations (69) and (70), we obtain $\alpha^2 + (3\lambda_1\lambda_2 - \lambda_2^2)\alpha = 0$, so when $\lambda_2 = 3\lambda_1$, we have $\alpha = 0$; when $\lambda_2 \neq 3\lambda_1$, we have $\alpha = 0$ or $\alpha = \lambda_2^2 - 3\lambda_1\lambda_2 \neq 0$.

1) $\lambda_2 = 3\lambda_1$. Then $\alpha = 0$, by the equation (70), we get ϕ^ζ is a constant, then ψ is a constant, so by the equation $(67a)$, we have $\lambda_1^2 = 0$, which is a contradiction with $\lambda_1 \neq 0$.

2) $\lambda_2 \neq 3\lambda_1$. $1'$. $\alpha = \lambda_2^2 - 3\lambda_1\lambda_2 \neq 0$. By the equation (70), we get ϕ^ζ is a constant, then ψ is a constant, and $\phi = c$ is a constant. by the equation $(67a)$, we have $\alpha = 3\lambda_1^2 - 3\lambda_1\lambda_2$ and $\lambda_2^2 = 3\lambda_1^2$.

So we get $\underline{\lambda_2^2 = 3\lambda_1^2,\ \alpha = 3\lambda_1^2 - 3\lambda_1\lambda_2 = \lambda_2^2 - 3\lambda_1\lambda_2,\ \alpha_i = 0,\ \zeta \neq 0,\ \eta \neq 0,\ \phi = c.}$

$2'$. $\alpha = 0$. By the equation $(67a)$, we have

$$\psi'' - \lambda_2\psi' + (3\lambda_1\lambda_2 - 3\lambda_1^2)\frac{\eta}{\zeta^2}\psi = 0. \tag{71}$$

by the equation (70), we get $\phi^{\zeta} = c_0 e^{(3\lambda_1-\lambda_2)t}$, then $\phi = ce^{\frac{3\lambda_1-\lambda_2}{\zeta}t}$, $\psi = c_1 e^{(3\lambda_1-\lambda_2)\frac{\eta}{\zeta^2}t}$, by the equation (71), we get $\frac{\eta}{\zeta^2} = \frac{3\lambda_1^2-\lambda_2^2}{(3\lambda_1-\lambda_2)^2}$.

So we have $\lambda_2 \neq 3\lambda_1,\ \alpha = 0,\ \alpha_i = 0,\ \frac{\eta}{\zeta^2} = \frac{3\lambda_1^2-\lambda_2^2}{(3\lambda_1-\lambda_2)^2},\ \phi = ce^{\frac{3\lambda_1-\lambda_2}{\zeta}t}$.

According to above discussions, we get the following Theorem:

Theorem 4.20. *Let $M = I \times_{\phi^{p_1}} F_1 \times_{\phi^{p_2}} F_2 \cdots \times_{\phi^{p_m}} F_m$ be a generalized Kasner space-time for $p_i \neq p_j$ for $i, j \in \{1, 2, 3\}$ and $\dim F_1 = \dim F_2 = \dim F_3 = 1$, and $P = \frac{\partial}{\partial t}$. Then $(M, \overline{\nabla})$ is Einstein with the Einstein constant α if and only if one of the following conditions is satisfied:*

(1) $\lambda_2^2 = 3\lambda_1^2,\ \alpha = 3\lambda_1^2 - 3\lambda_1\lambda_2 = \lambda_2^2 - 3\lambda_1\lambda_2,\ \alpha_i = 0,\ \zeta = \eta = 0$;

(2) $\lambda_2^2 = 3\lambda_1^2,\ \alpha = 3\lambda_1^2 - 3\lambda_1\lambda_2 = \lambda_2^2 - 3\lambda_1\lambda_2,\ \alpha_i = 0,\ \eta \neq 0,\ \phi = c$;

(3) $\lambda_2 \neq 3\lambda_1,\ \alpha = 0,\ \alpha_i = 0,\ \frac{\eta}{\zeta^2} = \frac{3\lambda_1^2-\lambda_2^2}{(3\lambda_1-\lambda_2)^2},\ \phi = ce^{\frac{3\lambda_1-\lambda_2}{\zeta}t}$.

4.3.6 Type (III) generalized Kasner space-times with a quarter-symmetric connection with constant scalar curvature

By Proposition 4.17, we get

$$\overline{S} = -2\zeta\frac{\phi''}{\phi} - (\eta + \zeta^2 - 2\zeta)\frac{(\phi')^2}{\phi^2} + 3(\lambda_1 + \lambda_2)\zeta\frac{\phi'}{\phi} + 3(\lambda_1^2 + \lambda_2^2 - 4\lambda_1\lambda_2). \tag{72}$$

$\zeta = \eta = 0$. Then $p_1 = p_2 = p_3 = 0$, and $\overline{S} = 3(\lambda_1^2 + \lambda_2^2 - 4\lambda_1\lambda_2)$.

$\zeta = 0,\ \eta \neq 0$. Then $[(ln\phi)']^2 = \frac{3(\lambda_1^2+\lambda_2^2-4\lambda_1\lambda_2)-\overline{S}}{\eta}$, so we have:

1) $\overline{S} > 3(\lambda_1^2 + \lambda_2^2 - 4\lambda_1\lambda_2)$, we have no solution.

2) $\overline{S} = 3(\lambda_1^2 + \lambda_2^2 - 4\lambda_1\lambda_2)$, then $\phi = c$.

3) $\overline{S} < 3(\lambda_1^2 + \lambda_2^2 - 4\lambda_1\lambda_2)$, then $\phi = c_0 e^{\pm\sqrt{\frac{3(\lambda_1^2+\lambda_2^2-4\lambda_1\lambda_2)-\overline{S}}{\eta}}t}$.

$\zeta \neq 0$, then $\eta \neq 0$. Putting $\phi = \psi^{\frac{2\zeta}{\eta+\zeta^2}}$, then

$$-\frac{4\zeta^2}{\eta+\zeta^2}\psi'' + \frac{6(\lambda_1+\lambda_2)\zeta^2}{\eta+\zeta^2}\psi' + (3\lambda_1^2 + 3\lambda_2^2 - 12\lambda_1\lambda_2 - \overline{S})\psi = 0. \tag{73}$$

So we get

1) $\overline{S} < \frac{9\zeta^2(\lambda_1+\lambda_2)^2}{4(\eta+\zeta^2)} + 3\lambda_1^2 + 3\lambda_2^2 - 12\lambda_1\lambda_2$,

$$\begin{aligned}\psi &= c_1 e^{\frac{\frac{3}{2}(\lambda_1+\lambda_2)+\sqrt{\frac{9(\lambda_1+\lambda_2)^2}{4} - \frac{(\overline{S}-3\lambda_1^2-3\lambda_2^2+12\lambda_1\lambda_2)(\eta+\zeta^2)}{\zeta^2}}}{2}t} \\ &+ c_2 e^{\frac{\frac{3}{2}(\lambda_1+\lambda_2)-\sqrt{\frac{9(\lambda_1+\lambda_2)^2}{4} - \frac{(\overline{S}-3\lambda_1^2-3\lambda_2^2+12\lambda_1\lambda_2)(\eta+\zeta^2)}{\zeta^2}}}{2}t};\end{aligned}$$

2) $\overline{S}=\frac{9\zeta^2(\lambda_1+\lambda_2)^2}{4(\eta+\zeta^2)}+3\lambda_1^2+3\lambda_2^2-12\lambda_1\lambda_2,\ \psi=c_1e^{\frac{3(\lambda_1+\lambda_2)}{4}t}+c_2te^{\frac{3(\lambda_1+\lambda_2)}{4}t}$;
3) $\overline{S}>\frac{9\zeta^2(\lambda_1+\lambda_2)^2}{4(\eta+\zeta^2)}+3\lambda_1^2+3\lambda_2^2-12\lambda_1\lambda_2$,

$$\begin{aligned}\psi &= c_1e^{\frac{3}{4}(\lambda_1+\lambda_2)t}cos\frac{\sqrt{\frac{(\overline{S}-3\lambda_1^2-3\lambda_2^2+12\lambda_1\lambda_2)(\eta+\zeta^2)}{\zeta^2}-\frac{9(\lambda_1+\lambda_2)^2}{4}}}{2}t\\ &+ c_2e^{\frac{3}{4}(\lambda_1+\lambda_2)t}sin\frac{\sqrt{\frac{(\overline{S}-3\lambda_1^2-3\lambda_2^2+12\lambda_1\lambda_2)(\eta+\zeta^2)}{\zeta^2}-\frac{9(\lambda_1+\lambda_2)^2}{4}}}{2}t\end{aligned}$$

According to above discussions, we get the following Theorem:

Theorem 4.21. *Let $M=I\times_{\phi^{p_1}}F_1\times_{\phi^{p_2}}F_2\cdots\times_{\phi^{p_m}}F_m$ be a generalized Kasner spacetime and $\dim F_1=\dim F_2=\dim F_3=1$, and $P=\frac{\partial}{\partial t}$. Then $(M,\overline{\nabla})$ has constant scalar curvature $\overline{S}$ if and only if one of the following conditions is satisfied:*
(1) $\zeta=\eta=0,\ \overline{S}=3(\lambda_1^2+\lambda_2^2-4\lambda_1\lambda_2)$.
(2) $\zeta=0,\ \eta\neq0$, *when* $\overline{S}>3(\lambda_1^2+\lambda_2^2-4\lambda_1\lambda_2)$, *we have no solution; when* $\overline{S}=3(\lambda_1^2+\lambda_2^2-4\lambda_1\lambda_2)$, *then* $\phi=c$; *when* $\overline{S}<3(\lambda_1^2+\lambda_2^2-4\lambda_1\lambda_2)$, *then* $\phi=c_0e^{\pm\sqrt{\frac{3(\lambda_1^2+\lambda_2^2-4\lambda_1\lambda_2)-\overline{S}}{\eta}}t}$.
(3) $\zeta\neq0$,
(a) $\overline{S}<\frac{9\zeta^2(\lambda_1+\lambda_2)^2}{4(\eta+\zeta^2)}+3\lambda_1^2+3\lambda_2^2-12\lambda_1\lambda_2$,

$$\begin{aligned}\psi &= c_1e^{\frac{\frac{3}{2}(\lambda_1+\lambda_2)+\sqrt{\frac{9(\lambda_1+\lambda_2)^2}{4}-\frac{(\overline{S}-3\lambda_1^2-3\lambda_2^2+12\lambda_1\lambda_2)(\eta+\zeta^2)}{\zeta^2}}}{2}t}\\ &+ c_2e^{\frac{\frac{3}{2}(\lambda_1+\lambda_2)-\sqrt{\frac{9(\lambda_1+\lambda_2)^2}{4}-\frac{(\overline{S}-3\lambda_1^2-3\lambda_2^2+12\lambda_1\lambda_2)(\eta+\zeta^2)}{\zeta^2}}}{2}t};\end{aligned}$$

(b) $\overline{S}=\frac{9\zeta^2(\lambda_1+\lambda_2)^2}{4(\eta+\zeta^2)}+3\lambda_1^2+3\lambda_2^2-12\lambda_1\lambda_2,\ \psi=c_1e^{\frac{3(\lambda_1+\lambda_2)}{4}t}+c_2te^{\frac{3(\lambda_1+\lambda_2)}{4}t}$;
(c) $\overline{S}>\frac{9\zeta^2(\lambda_1+\lambda_2)^2}{4(\eta+\zeta^2)}+3\lambda_1^2+3\lambda_2^2-12\lambda_1\lambda_2$,

$$\begin{aligned}\psi &= c_1e^{\frac{3}{4}(\lambda_1+\lambda_2)t}cos\frac{\sqrt{\frac{(\overline{S}-3\lambda_1^2-3\lambda_2^2+12\lambda_1\lambda_2)(\eta+\zeta^2)}{\zeta^2}-\frac{9(\lambda_1+\lambda_2)^2}{4}}}{2}t\\ &+ c_2e^{\frac{3}{4}(\lambda_1+\lambda_2)t}sin\frac{\sqrt{\frac{(\overline{S}-3\lambda_1^2-3\lambda_2^2+12\lambda_1\lambda_2)(\eta+\zeta^2)}{\zeta^2}-\frac{9(\lambda_1+\lambda_2)^2}{4}}}{2}t\end{aligned}$$

Remark 13. *When $\lambda_1=\lambda_2=1$, we get Propositions 32, 33 Theorems 35, 37, 36, in* [12] *by Propositions 4.16, 4.17, Theorems 4.19 − 4.21, respectively.*

Acknowledgement. This work was supported by NSFC No.11271062 and NCET-13-0721. We would like to thank the referee for his(her) careful reading and helpful comments.

References

[1] R. Bishop, B. O'Neill, Manifolds of negative curvature, Trans. Am. Math. Soc.145(1969) 1-49.

[2] F. Dobarro, E. Dozo, Scalar curvature and warped products of Riemannian manifolds, Trans. Am. Math. Soc. 303(1987) 161-168.

[3] P. Ehrlich, Y. Jung, S. Kim, Constant scalar curvatures on warped product manifolds, Tsukuba J. Math. 20(1996) No.1 239-265.

[4] L. Alías, A. Romero, M. S'anchez, Spacelike hypersurfaces of constant mean curvature and Clabi-Bernstein type problems, Tohoku Math. J. 49(1997) 337-345.

[5] M. Fernández-López, E. García-Río, D. Kupeli, B. Ünal, A curvature condition for a twisted product to be a warped product, Manu. math. 106(2001), 213-217.

[6] H. A. Hayden, Subspace of a space with torsion, Proc. Lond. Math. Soc. 34(1932) 27-50.

[7] K. Yano, On semi-symmetric metric connection, Rev. Roumaine Math. Pures Appl. 15(1970) 1579-1586.

[8] Golab, S., On semi-symmetric and quarter-symmetric linear connections, Tensor N.S., 29(1975), 249-254.

[9] F. Dobarro, B. Ünal, Curvature of multiply warped products, J. Geom. Phys. 55(2005) 75-106.

[10] S. Sular and C. Özgür, Warped products with a semi-symmetric metric connection, Taiwanese J. Math. 15 (2011), no. 4, 1701C1719.

[11] S.Sular and C.Özgür, Warped products with a semi-symmetric non-metric connection, Arab. J. Sci. Eng. 36 (2011), no. 3, 461C473.

[12] Y. Wang, Multiply Warped products with a semi-symmetric metric connection, Abstract and Applied Analysis. Vol 2014, Article ID 742371, 12pages.

[13] Y. Wang, Curvature of multiply warped products with an affine connection, Bull. Korean Math. Soc. 50 (2013), No. 5, pp. 1567C1586

[14] M. Tripathi, A new connection in a Riemannian manifold, arXiv:1207.0199